

**СООТВЕТСТВУЕТ  
ГОСТ 7.56-2002**  
ПЕЧАТНОЕ ИЗДАНИЕ  
ISSN 2312-8089

№ 4 (107). Ч.1. ФЕВРАЛЬ 2021

# ВЕСТНИК НАУКИ И ОБРАЗОВАНИЯ

НАУЧНО-МЕТОДИЧЕСКИЙ ЖУРНАЛ

 **РОСКОМНАДЗОР**

ПИ № ФС 77-50633 • Эл № ФС 77-58456

НАУЧНО-МЕТОДИЧЕСКИЙ ЖУРНАЛ «ВЕСТНИК НАУКИ И ОБРАЗОВАНИЯ» № 4 (107) Ч.1. 2021



ИЗДАТЕЛЬСТВО «ПРОБЛЕМЫ НАУКИ»

[HTTPS://SCIENCEPROBLEMS.RU](https://scienceproblems.ru)

ЖУРНАЛ: [HTTP://SCIENTIFICJOURNAL.RU](http://scientificjournal.ru)

 НАУЧНАЯ ЭЛЕКТРОННАЯ  
БИБЛИОТЕКА  
**LIBRARY.RU**



9 772312 808001

**ВЕСТНИК НАУКИ  
И ОБРАЗОВАНИЯ**

2021. № 4 (107). Часть 1



Москва  
2021

# Содержание

<b>ТЕХНИЧЕСКИЕ НАУКИ</b> .....	<b>6</b>
<i>Махмудова Н.Р.</i> АНАЛИЗ ОБЛАЧНЫХ ТЕХНОЛОГИЙ И ВЗАИМОДЕЙСТВИЯ ТЕЛЕКОММУНИКАЦИОННЫХ СИСТЕМ И СЕТЕЙ / <i>Makhmudova N.R.</i> ANALYSIS OF CLOUD TECHNOLOGIES AND INTERACTIONS OF TELECOMMUNICATION SYSTEMS AND NETWORKS .....	6
<b>ФИЛОЛОГИЧЕСКИЕ НАУКИ</b> .....	<b>11</b>
<i>Сайдазимова У.Т.</i> КОРЕЙСКАЯ КЛАССИЧЕСКАЯ ПРОЗА XVII – XVIII ВЕКОВ: ОСОБЕННОСТИ ЕЕ СТАНОВЛЕНИЯ И РАЗВИТИЯ / <i>Saydazimova U.T.</i> KOREAN CLASSICAL PROSE OF THE XVII-XVIII CENTURIES, THE FEATURES OF ITS FORMATION AND DEVELOPMENT .....	11
<i>Хакимова Д.</i> ТЕМАТИЧЕСКИЕ ГРУППЫ И МОТИВАЦИОННЫЕ ОСОБЕННОСТИ ГОРНОЙ ЛЕКСИКИ ПРИ ОБУЧЕНИИ ТЕРМИНОЛОГИИ ГОРНОГО ДЕЛА / <i>Khakimova D.</i> THEME GROUPS AND MOTIVATIVE FEATURES OF MINING VOCABULARY IN TEACHING THE TERMINOLOGY OF MINING .....	25
<b>ЮРИДИЧЕСКИЕ НАУКИ</b> .....	<b>28</b>
<i>Бояринцева И.А.</i> МОДЕЛИ МЕСТНОГО САМОУПРАВЛЕНИЯ В ИМПЕРАТОРСКОЙ РОССИИ XIX ВЕКА И СОВРЕМЕННЫЕ ПАРАЛЛЕЛИ / <i>Boyarintseva I.A.</i> MODELS OF LOCAL SELF- GOVERNMENT IN IMPERIAL RUSSIA OF THE XIX CENTURY AND MODERN PARALLELS.....	28
<i>Гладкая Ю.Д.</i> ПРАВО МЕТАМОДЕРНА В УСЛОВИЯХ ГЛОБАЛИЗАЦИИ / <i>Gladkaya Yu.D.</i> THE RIGHT OF METAMODERN IN THE CONDITIONS OF GLOBALIZATION .....	31
<i>Копейкина И.В.</i> КРИТЕРИИ ДОПУСТИМОСТИ ДОКАЗАТЕЛЬСТВ В УГОЛОВНОМ ПРОЦЕССЕ / <i>Kopeikina I.V.</i> CRITERIA FOR THE ADMISSIBILITY OF EVIDENCE IN CRIMINAL PROCEEDINGS .....	34
<i>Лучина А.Ю.</i> ФОРМЫ МНОЖЕСТВЕННОСТИ ПРЕСТУПЛЕНИЙ ПО УГОЛОВНОМУ ПРАВУ РОССИЙСКОЙ ФЕДЕРАЦИИ И АВСТРИЙСКОЙ РЕСПУБЛИКИ / <i>Luchina A.Yu.</i> FORMS OF MULTIPLE CRIMES UNDER THE CRIMINAL LAW OF THE RUSSIAN FEDERATION AND THE REPUBLIC OF AUSTRIA.....	37
<b>ПЕДАГОГИЧЕСКИЕ НАУКИ</b> .....	<b>42</b>
<i>Амерова Р.Х., Глотова Л.Р.</i> РОЛЬ ЛИНГВИСТИЧЕСКИХ СЛОВАРЕЙ ПРИ ОБУЧЕНИИ СТУДЕНТОВ-ИНОСТРАНЦЕВ РУССКОМУ ЯЗЫКУ / <i>Amerova R.H., Glotova L.R.</i> ROLE OF LINGUISTIC DICTIONARIES IN TEACHING FOREIGN STUDENTS IN RUSSIAN .....	42
<b>МЕДИЦИНСКИЕ НАУКИ</b> .....	<b>45</b>
<i>Каюмов У.К., Бекматова Ш.К., Рахманова С.С., Саипова М.Л.</i> ОЦЕНКА СОСТОЯНИЯ СВОЕГО ЗДОРОВЬЯ СРЕДИ ЛИЦ С ИЗБЫТОЧНОЙ МАССОЙ ТЕЛА В РАЗЛИЧНЫЕ ВОЗРАСТНЫЕ ПЕРИОДЫ / <i>Kayumov</i>	

<i>U.K., Bekmatova Sh.K., Rakhmanova S.S., Saipova M.L.</i> ASSESSMENT OF THE STATE OF THEIR HEALTH AMONG OVERWEIGHT PEOPLE IN DIFFERENT AGE PERIODS .....	45
<i>Хатамова Д.Т., Матмуратова С.О., Рахманова С.С., Зиямухамедова М.М.</i> ОЦЕНКА СОСТОЯНИЯ МАССЫ ТЕЛА У ЖЕНЩИН РЕПРОДУКТИВНОГО ВОЗРАСТА / <i>Khatamova D.T., Matmuratova S.O., Rakhmanova S.S., Ziyamukhamedova M.M.</i> ASSESSMENT OF THE STATE OF BODY WEIGHT IN WOMEN OF REPRODUCTIVE AGE .....	50
<i>Набиева Ф.С., Душанова Г.А., Бобокулов О.О.</i> ЗНАЧЕНИЕ ИММУНОФЕРМЕНТНОГО АНАЛИЗА В ДИАГНОСТИКЕ ИНФЕКЦИОННЫХ ЗАБОЛЕВАНИЙ / <i>Nabieva F.S., Dushanova G.A., Bobokulov O.O.</i> THE IMPORTANCE OF IMMUNO-ENZYMAL ANALYSIS IN THE DIAGNOSIS OF INFECTIOUS DISEASES .....	54
<i>Sadikov N., Xu Chao Yue, Odilov B., Xin Zhi Hong, Zhang Zhao Hua.</i> THE EFFECTIVENESS OF USING PREDNISOLONE IN CHILDREN WITH COMMUNITY–ACQUIRED PNEUMONIA / <i>Садиков Н., Шу Чао Юе, Одиллов Б., Шин Жи Хонг, Жанг Жао Хуа.</i> ЭФФЕКТИВНОСТЬ ПРЕДНИЗОЛОНА У ДЕТЕЙ С ВНЕБОЛЬНИЧНОЙ ПНЕВМОНИЕЙ.....	57
<i>Мирзаева Д.А., Худайкулова Ф.Х.</i> ОПРЕДЕЛЕНИЕ ТЯЖЕСТИ ОСТЕОАРТРОЗА У БОЛЬНЫХ С ОЖИРЕНИЕМ / <i>Mirzaeva D.A., Khudaykulova F.Kh.</i> DETERMINATION OF THE SEVERITY OF OSTEOARTHRISIS IN PATIENTS WITH OBESITY.....	66
<i>Ибрагимова Н.С., Ибрагимов Б.Ф., Махматкулов Р.А.</i> ДИАГНОСТИЧЕСКИЕ КРИТЕРИИ СИНДРОМА ПОЛИКИСТОЗНЫХ ЯИЧНИКОВ / <i>Ibragimova N.S., Ibragimov B.F., Makhmatkulov R.A.</i> DIAGNOSTIC CRITERIA FOR POLYCYSTIC OVARY SYNDROME .....	70
<i>Азимов С.И., Гуламов М.Х., Каримов Б.С.</i> ФУНКЦИОНАЛЬНОЕ СОСТОЯНИЕ И КАЧЕСТВО ЖИЗНИ У ПАЦИЕНТОВ С ДОБРОКАЧЕСТВЕННОЙ ГИПЕРПЛАЗИЕЙ ПРЕДСТАТЕЛЬНОЙ ЖЕЛЕЗЫ ПОСЛЕ ЭМБОЛИЗАЦИИ АРТЕРИЙ ПРОСТАТЫ / <i>Azimov S.I., Gulatov M.Kh., Karimov B.S.</i> FUNCTIONAL STATE AND QUALITY OF LIFE IN PATIENTS WITH BENIGN PROSTATIC HYPERPLASIA AFTER PROSTATIC ARTERY EMBOLIZATION .....	73
<i>Rustamova N.H., Vobojonova Sh.D.</i> INDICATORS OF BLOOD FERRITIN LEVEL IN PATIENTS WITH APLASTIC ANEMIA / <i>Рустамова Н.Х., Вобожонова Ш.Д.</i> ПОКАЗАТЕЛИ УРОВНЯ ФЕРРИТИНА КРОВИ У БОЛЬНЫХ С АПЛАСТИЧЕСКОЙ АНЕМИЕЙ .....	77
<b>ИСКУССТВОВЕДЕНИЕ .....</b>	<b>81</b>
<i>Плужникова Д.В.</i> НОВАЦИИ В БАЛЕТАХ М. ФОКИНА «ПЕТРУШКА» И В. НИЖИНСКОГО «ВЕСНА СВЯЩЕННАЯ» КАК НАЧАЛО ИНОГО ПУТИ РАЗВИТИЯ БАЛЕТА XIX ВЕКА / <i>Pluzhnikova D.V.</i> INNOVATIONS IN THE BALLETS OF M. FOKIN "PETRUSHKA" AND V. NIJINSKY "SACRED SPRING" AS THE BEGINNING OF A DIFFERENT PATH OF THE BALLET DEVELOPMENT IN THE XIX CENTURY .....	81

<b>ПСИХОЛОГИЧЕСКИЕ НАУКИ.....</b>	<b>84</b>
<i>Созина М.В.</i> ПРОФИЛАКТИКА ПРОФЕССИОНАЛЬНЫХ КОНФЛИКТОВ В УЧРЕЖДЕНИЯХ СОЦИАЛЬНОЙ ЗАЩИТЫ НАСЕЛЕНИЯ САНКТ-ПЕТЕРБУРГА / <i>Sozinova M.V.</i> PREVENTION OF PROFESSIONAL CONFLICTS IN SOCIAL PROTECTION INSTITUTIONS OF THE POPULATION OF ST. PETERSBURG.....	84
<b>ПОЛИТИЧЕСКИЕ НАУКИ.....</b>	<b>90</b>
<i>Волужев Е.А., Саяпина О.Н.</i> АНТРОПОГЕННЫЕ ПРОБЛЕМЫ ПРОФИЛАКТИКИ ЛЕСНЫХ ПОЖАРОВ НА ТЕРРИТОРИИ ХАБАРОВСКОГО КРАЯ / <i>Voluzhev E.A., Sayapina O.N.</i> ANTHROPOGENIC PROBLEMS OF FOREST FIRE PREVENTION IN THE Khabarovsk REGION .....	90
<i>Sagdullayeva D.S.</i> THE ROLE OF THE MASS MEDIA IN THE DEVELOPMENT OF EMOTIONAL INTELLIGENCE, THE LEADERS OF THE YOUTH / <i>Сагдуллаева Д.С.</i> РОЛЬ СРЕДСТВ МАССОВОЙ ИНФОРМАЦИИ В РАЗВИТИИ ЭМОЦИОНАЛЬНОГО ИНТЕЛЛЕКТА ЛИДЕРОВ МОЛОДЕЖИ .....	95

# INDICATORS OF BLOOD FERRITIN LEVEL IN PATIENTS WITH APLASTIC ANEMIA

Rustamova N.H.<sup>1</sup>, Bobojonova Sh.D.<sup>2</sup>  
Email: Rustamova6107@scientifictext.ru

<sup>1</sup>Rustamova Nilufar Hurmatovna – Assistant,  
DEPARTMENT OF FACULTY AND HOSPITAL THERAPY,  
URGENCH BRANCH

TASHKENT MEDICAL ACADEMY, URGENCH;  
<sup>2</sup>Bobojonova Shohista Davronbekovna – Assistant,  
DEPARTMENT OF HEMATOLOGY, TRANSFUSIOLOGY AND LABORATORY SCIENCE,  
TASHKENT MEDICAL ACADEMY, TASHKENT,  
REPUBLIC OF UZBEKISTAN

**Abstract:** iron overload develops as a result of its excessive accumulation in parenchymal organs, which subsequently leads to their damage and dysfunction. The assessment of the body's iron storage is carried out indirectly by determining the concentration of ferritin in the blood serum. However, the concentration of ferritin in the patient's blood serum is associated with the presence of inflammation, with ascorbate deficiency, and also depends on liver function, which significantly limits the diagnostic value. Iron by nature is a ferromagnet, its nuclei have their own magnetic moments, which, when hitting a strong magnetic moment, line up parallel to each other and greatly shorten the relaxation time of the magnetic field. In the last 20 years, magnetic resonance imaging (MRI) has been actively used to determine and calculate the amount of accumulated iron in various organs - liver, heart, pancreas, pituitary gland, etc.

**Keywords:** aplastic anemia, myelodysplastic syndrome, magnetic resonance imaging, ferritin.

## ПОКАЗАТЕЛИ УРОВНЯ ФЕРРИТИНА КРОВИ У БОЛЬНЫХ С АПЛАСТИЧЕСКОЙ АНЕМИЕЙ Рустамова Н.Х.<sup>1</sup>, Бобожонова Ш.Д.<sup>2</sup>

<sup>1</sup>Рустамова Нилуфар Хурматовна – ассистент,  
кафедра факультетской и госпитальной терапии,  
Ургенчский филиал

Ташкентская медицинская академия, г. Ургенч;  
<sup>2</sup>Bobojonova Shohista Davronbekovna – ассистент,  
кафедра гематологии, трансфузиологии и лабораторного дела,  
Ташкентская медицинская академия, г. Ташкент,  
Республика Узбекистан

**Аннотация:** перегрузка железом развивается в результате избыточного его накопления в паренхиматозных органах, впоследствии приводит к их повреждению и дисфункции. Оценку запаса железа в организме проводят косвенно - путем определения концентрации ферритина в сыворотке крови. Однако концентрация ферритина в сыворотке крови больного связана с наличием воспаления, с дефицитом аскорбата, зависит также от функции печени, что значительно ограничивает диагностическое значение. Железо по своей природе - ферромагнетик, его ядра обладают собственными магнитными моментами, которые при попадании в сильное магнитное поле выстраиваются параллельно друг другу и сильно укорачивают время релаксации магнитного поля. В последние 20 лет магнитно-резонансную томографию (МРТ) активно используют для определения и подсчета количества накопленного железа в различных органах - печени, сердце, поджелудочной железе, гипофизе и др.

**Ключевые слова:** апластическая анемия, миелодиспластическим синдром, магнитно-резонансная томография, ферритин.

UDC 616.155.194.7:612.398-07

A necessary component of modern therapy for aplastic anemia (AA) is an adequate transfusion supply, in particular, transfusion of erythrocyte mass [1, 2]. At the same time, it is known that transfusions of donor erythrocytes cause the development of secondary siderosis, and the carriage of hereditary hemochromatosis gene mutations aggravates the formation of iron overload in the body [3, 4].

In patients with chronic hepatitis and cirrhosis of the liver, as a result of repeated blood transfusions, iron overload may also develop, which exacerbates the risk of progression of chronic liver disease. In the research of M.L. Zubkina et al. Showed that antiviral therapy in combination with massive bloodletting, clinical observation of a patient with chronic hepatitis B allowed stabilization of chronic liver disease [5].

In the studies of V.G. Savchenko. et al. have shown the results of chelation therapy in transfusion-dependent patients with myelodysplastic syndrome (MDS), whose serum ferritin values were 1000  $\mu\text{g} / \text{L}$  or more. In a prospective open multicenter study, the safety and efficacy of deferasirox therapy in transfusion-dependent patients with MDS, as well as rare forms of anemia, was evaluated. The results showed that chelation therapy can effectively reduce the iron content in the patient's body. The mean serum ferritin value decreased from 3837.2 to 2269.2  $\mu\text{g} / \text{L}$ . What is a statistically significant criterion for the effectiveness of chelation therapy [6].

**Materials and research methods.** The object of the study was 20 patients with a diagnosis of AA, who were treated at the Research Institute of Hematology and Blood Transfusion clinic. The age of the patients was from 19 to 67 years old, on average  $39.79 \pm 3.64$  years. 11 women, 9 men, 12 urban residents, 8 rural residents. All patients underwent a general blood test on a MINDRAYBC-2300 hematology analyzer, ferritin values were determined on a RENDEX biochemical analyzer (Daytona). The ferritin concentration norm was 10-120 ng / L. Myelogram and coagulogram were calculated. The anamnesis was studied, in particular the volume of transfused blood for the entire period of the disease, which ranged from 1518 liters to 9960 liters, the average volume was  $5778.11 \pm 633.87$  liters. The duration of the disease was 1-12 years, on average  $2.86 \pm 0.64$  years.

**Results and discussion.** Blood transfusions are widely used in clinical practice as a quick and effective method for correcting anemia. Most often, transfusions are used in hematology oncology, including in patients with aplastic anemia. Indications for blood transfusion occurs when hemoglobin decreases below 65-80 g / l, if it is accompanied by clinical symptoms of hemodynamic disturbance (tachycardia, shortness of breath, decreased blood pressure, etc.). However, existing clinical guidelines for the treatment of anemia do not recommend long-term blood transfusion therapy in hematological cancer patients due to the risk of iron overload and sensitization to blood antigens. Patients with AA deserve special attention, who are forced to be on constant blood transfusions for life, and therefore have a high risk of iron overload. In our study, we observed 20 patients with AA who received regular blood transfusions (once a month). The results of the study are shown in table 1.

Table 1. Indicators of blood ferritin concentration depending on some indicators

№	Indicators	Ferritin indices ng/l M ± m	Reliability
1	Age		
a	19-40 year (n=11)	1155±145,2*	(P>0,5)
B	41-67 year (n=9)	884±139,1*	
2	Blood transfusion volume		
a	1-6 liter (n=11)	801±128,4**	(P<0,05)
B	7-9 liter (n=9)	1317±140,5**	
3	Duration of illness		
a	6 month- 1 year 11 month (n=12)	693,6±120,1**	(P.<0,05)
B	2-12 year (n=8)	1260±145,5**	
4	Sex		
a	Men (n=9)	985±119,5*	(P>0,5)
B	Women (n=11)	1073±123,2*	

Note:

\* - the difference is insignificant (P> 0.5)

\*\* - the difference is significant (P <0.05)

The study of blood ferritin indicators depending on age indicates that at the age of 19-40 years, the concentration of ferritin was  $1155 \pm 145.2$  ng / l, at the age of 41-67 it was  $884 \pm 139.1$  ng / l, however, there was no significant difference in indicators revealed (P> 0.5). The study of ferritin concentration depending on the volume of blood transfusion made it possible to establish a significant difference in indicators: in patients who received blood transfusion in the amount of 1-6 liters per year, ferritin indicators were  $801 \pm 128.4$  ng / l, and in patients who received more than 6-9 liters of weight per year the level of ferritin was  $1317 \pm 140.5$  ng / l, the difference is significant (P <0.05).

Consequently, the greater the volume of blood transfusion, the higher the risk of developing iron overload.

Ferritin indices were also studied depending on the duration of the disease and the following was found: with the duration of the disease up to 2 years ago, the ferritin level was  $693.6 \pm 120.1$  ng / l, and with the duration of the disease 2-12 years  $1260 \pm 145.5$  ng / l, the difference is significant (P. <0.05). Thus, the risk of developing iron overload directly depends on the duration of the disease: the longer the duration of the disease, the higher the ferritin values. In our studies, the dependence of ferritin indicators on gender was studied, while the data showed no relationship (P> 0.5).

In the literature we studied, there are very few data on the study of iron overload in transfusion-dependent hematological patients, including patients with AA. Retrospective studies of the medical history of patients with AA show that practitioners do not pay due attention to the issues of iron overload, ferritin indicators are not studied, and therefore timely chelation therapy is not carried out. As literature sources show (1, 2, 3, 4, 5), iron overload causes dysfunction of the internal organs - heart, liver, spleen, pancreas, pituitary gland, etc. heart, spleen, etc. When iron overload syndrome is established, pathogenetic therapy (chelation) is necessary.

### Conclusions:

1. In patients with AA, with a disease duration of 1 year or more, in 100% of cases, ferritin indices increase by almost 10 times.

2. Ferritin values in patients with AA depend on the duration of the disease and on the volume of blood transfusions received: the longer the duration of the disease and the volume of blood transfused, the higher the ferritin values and the higher the risk of hemosiderosis.

3. Indicators of ferritin does not depend on the sex and age of the patients.



### *References / Список литературы*

1. *Savchenko V.G., Abdulkadyrov K.M., Maschan A.A. et al.* An open multicenter study of deferasirox in the treatment of post-transfusion iron overload in patients with myelodysplastic syndromes, thalassemia, and other forms of anemia. *Ros, journal Hematology and transfusiology*. 2015. vol. 60, no. 4 (7-13).
2. *Powell L.W.* Hereditary hemochromatosis and iron overload diseases. *J. Gastroenterol Hepatol*, 2002; 17: 191-5.
3. *Angelucci E., Brittenham G.M., McLaren C E., Ripalti M., Baronciani D., Giardini C. et al.* Hepatic iron concentration and total body iron stores in thalassemia major. *N Engl J Med.*, 2000; 343 (5): 327-31.
4. *Bacon B.R., Adams P.S., Kowdley K.V., Powell L.W., Tavill A.S.* Diagnosis and management of hemochromatosis: 2011 Practice Guideline by the American Association for the Study of Liver Diseases. *Hepatology*, 2011; 54: 328-43.
5. *Tsvetaeva N.V., Levina A.A.* Kinetics of iron excretion by desferal and plasmapheresis in the treatment of patients with secondary hemochromatosis. In the book: III All-Union Congress of Hematologists. M., 1991; h. 2: 245-246.
6. *Niederrau C.* Hereditary Hemochromatosis. *Med. Klin.*, 2009; 104 (12): 931-946.