

ҚОН ҚҮЙИШ ОРҚАЛИ ЮҚАДИГАН БАКТЕРИАЛ ИНФЕКЦИЯЛАР

Ш. Д. Бобожонова

Тошкент тиббиёт академияси асистенти

Л. Ж. Қурбонова

РИГИАТМси клиник диагностик лаборатория шифокори

Н. Х. Рустамова

Тошкент тиббиёт академияси Урганч филиали асистенти

АННОТАЦИЯ

Трансфузиологияда юқадиган инфекциялар юқиши хавфи бугунги кунда ҳар қачонгидан пастроқ бўлсада, хавфсиз қон махсулотларини етказиб беришда, маълум бўлган ва хали аниқланмаган патогенлар билан ифлосланиш остида қолмоқда. Фақатгина донорларни танлашни такомиллаштириш ва амалга ошириш, сезгир скрининг синовлари ва самарали инактивация протседуралари трансфузион юқадиган инфекцияларни юқиши хавфини йўқ қилишни ёки хеч бўлмагандан камайтиришни таъминлаш мумкин. Бундан ташқари, қон қўйилиши орқали юқиши мумкин бўлган юқимли касалликларга оид доимий маълумот ва замонавий маълумотлар қон қўйиш орқали юқадиган касалликларни назорат қилишининг мухим таркибий қисми бўлган нохуш ходисалар тўғрисида хабар бериш учун зарурдир. Шундай қилиб, трансфузион тиббиёт билан шуғулланадиган барча тамонларнинг шу жумладан миллий қон назорати тизимларининг хамкорлиги, маълум бўлган ва пайдо бўлаётган (аниқланмаган) қон билан юқадиган патогенлардан хавфсиз қон махсулоти таъминотини химоя қилиш учун жуда мухим.

Калит сўзлар: Трансфузиология, донор, бактерия, тромбоцитлар концентрати, плазма, гемокомпонент, қизил қон таркибий қисмлари, инактивация.

КИРИШ

Тиббиёт тарихида bemорларни одам ёки хайвон қони билан даволашга уринишларни тавсифловчи дастлабки хабарлар мавжуд бўлсада, трансфузион тиббиёт бу ўтган асрнинг иккинчи ярмидан бошлаб ривожланган нисбатан ёш соҳадир. Шу билан бирга, жуда тезкор равишда, ушбу терапевтик ёндашувлар

донорлар ва қабул қилувчилар ўртасида қизил қон танаачалари ва плазмасининг мувофиқлиги ва юқумли касалликларни юқтириш имконияти каби муоммоларини ўз ичига олганлиги аниқ бўлди. [1, 2]. Илгари, қон қўйиш оралиқли юқадиган юқумли касалликлар bemорлар ва шифокорлар тамонидан муқаррар деб хисобланган бўлса, бугунги кунда кам хавфли қон таъминоти кутилмоқда. Бугунги кунда донорларни баҳолаш, лаборатория текширувлари ва патогенларни инактивация қилиш муолажалари трансфузион трансмиссив инфекциялар хавфини камайтириш учун хал қилувчи восита хисобланади, аммо барча хавфларни тўлиқ бартараф этмайди. Шу билан бирга, ушбу ютуқлар трансфузион тиббиётни тобора хавфсизроқ маҳсулотларга, харажатларни барқарор равишда ошириб боришига олиб келди . Амалдаги харакатлар ва стратегиялар қон қўйиш билан боғлиқ хавфларини камайтиришга катта ёрдам берди. Дарҳақиқат, бугунги кунда ифлосланган қон маҳсулотлари тамонидан инфекцияларни юқтириш хавфи ўттиз йил аввалги кўрсаткичга қараганда анча паст. Қон таъминотининг яхлитлигини, тозалигини ва этарлилигини сақлаб қолиш учун донорларни янги скрининг текширувлари, донорларнинг қон таркибий қисмлари патогеннинг инактивация қилиш жараёнлари, хаддан ташқари қатъий истисно қилиш мезонлари сабабли потенциал донорларнинг асоссиз йўқолишига қарши мувозанатлаштирилиши керак. Ушбу сай-харакатлар янги пайдо бўлаётган трансфузион трансмиссив инфекциялар тахдидларни аниқлашга ёрдам берадиган миллий ва халқаро хавфсизлик назорати тармоқлари тамонидан идеал тарзда қўллаб-қувватланади; сифатни таъминлаш, сифати ва трансфузион занжиридаги барча босқичларни кузатиш қобилиятини осонлаштириш орқали .[4-6].

АДАБИЁТЛАР ТАҲЛИЛИ ВА МЕТОДОЛОГИЯ

Бактериал инфекциялар хавфи трансфузия билан боғлиқ касалликлар ва ўлимнинг асосий сабаби бўлиб, қисман бошқа хавфларнинг камайиши билан боғлиқ [4, 7-10]. Бактерияларнинг кириш дарвозаси — терининг, овқат хазм қилиш аъзолари ва нафас йўлларининг шиллиқ қавати микрожарохати хисобланади. Контаминацияланган донор қонини қўйиш пайтида оғир бактериал асоратларни ривожланиш хавфи олдиндан маълум бўлган. Бироқ қон қўйишнинг бактериал хавфсизлиги муаммоси, айниқса, ўтган асрнинг 80-йиллари ўрталарида тромбоцитлар концентратидан (ТК) кенг фойдаланиш бошланиши муносабати билан жуда долзарб бўлиб қолди. Тромбоцитлар концентратларини қўйиш гемабластоз, апластик анемия билан оғриган bemорларда, шунингдек,

травмада кўп қон йўқотиши фонида ривожланадиган анемия ва тромбоцитопения билан оғриган беморларда тромбоцитопеник геморрагик синдром оқибатида келиб чиқадиган ўлим холатларини сезиларли даражада камайтириши мумкин. Донор тромбоцитларига талаб йил сайин ортиб бормоқда. Ушбу гемокомпонент, тайёрлаш усулидан қатъий назар, 20-22 ° С хароратда сақланади, худди шу харорат бактерияларнинг хаётий фаолияти учун мақбулдир. Бактериал ифлосланиш тромбоцитлар концентратларида қизил қон таркибий қисмларига қараганда тез-тез учрайди, чунки кўп микроорганизмлар одатда ТК (20-24 ° С) учун ишлатиладиган сақлаш шароитида яшashi ва тарқалиши мумкин, аммо қизил қон таркибий қисмлари учун камроқ (1-6) ° С) [8–11]. Мавжуд назорат қилиш усуллари хар доим хам қонда бактериялар мавжудлигини бутунлай чиқариб ташлашга имкон бермаслиги сабабли, уларни инактивациялаш усуллари кенг қўлланилади. Бироқ, ушбу усулларнинг ривожланиши катта қийинчиликлар билан боғлиқ. Хозирги вақтда плазма ва тромбоцит компонентларини инактивация қилиш тизимлари мавжуд. Эритроцит таркибий қисмлари учун бундай тизимлар хали мавжуд эмас . Шуни хам ёдда тутиш керакки, инактивация қилинган бактериялар ва уларнинг токсинларининг реципиент организмига таъсирини эмас, балки қон қўйишдан олдин сақлаш пайтида бактерияларни кўпайтириш имкониятини олдини олади. Бактериал ифлосланишни назорат қилишда инструментал усуллар билан бир қаторда оддий қўлда ишлатиладиган усуллардан фойдаланилади. Қон компонентларини тайёрлаш ва қўйишнинг мунтазам бактериал ифлосланишни сифатни назорат қилиш. [8–11]. Қон таркибий қисмларининг бактериал ифлосланишидан хабардорлик ва клиник долзарблигини ошириш натижасида илгари Американинг Қон Банки Ассоциацияси 2004 йилда бактериал трансфузион трансмиссив инфекцияларни камайтириш стандартларини чиқарди [12]. Хусусан, тромбоцитларни бактериал текшириш қон қўйиш хавфсизлигини яхшилаш бўйича самарали чоралар. Клиник оқибатларга олиб келадиган бактериал трансфузион трансмиссив инфекция билан касалланишнинг тахминий даражаси 70,000 дан 118,000 гача қўйилган тромбоцитар концентратлардан , асосан бактериялар турига ва уларнинг патогенлигига боғлиқ [13, 8, 14]. Қон компонентлари бактериал ифлосланишни аниқлаш бўйича Америка Қизил Хоч ташкилотининг хисоботи шуни кўрсатдики, тромбоцит концентратлари трансфузияси билан боғлиқ бўлган сепсис холатларида аниқланган организмлар қаторига (75% га яқини) граммусбат аэроб патогенлардир.

Бактерияларнинг ифлосланиш хавфини камайтириш бўйича чоратадбирлар трансфузион занжирининг турли босқичларига қаратилган ва уларнинг олти жихат бўйича таснифлаш мумкин:

1. Донорлик хуқуқи: Асимтоматик донор бактеремиясини камайтириш учун, яқин қунларда стоматологик муолажалар, кичик жаррохлик амалиёти ёки тана харорати кўтарилигдан донорликдан чиқариш керак.

2. Махсулотни мақбул қайта ишлаш ва сақлаш: масул ходимларни доимий ўқитиш ва назорат қилиш юқори сифат стандартлари ва хавфсиз махсулот учун асосий элементдир. Шунингдек, махсулотларнинг яхлитлигини таъминлаш учун доимий сақлаш хароратини (Эр.масса учун 4 ° C ва ТК учун 22-24 ° C) сақлаш керак.

3. Терини тайёрлаш: Донорлар қўлларини яхшилаб дезенфекциялаш, пункция жойида қолган бактериялар сонини камайтиришда хал қилувчи ахамиятга эга эканлиги исботланган [16-18].

4. Дастребаки тўлиқ қон тўпламини олиб ташлаш: Тўплам сумкасидан дастребаки 30-40 мл қонни чиқариб ташлаш тери бактерияларининг ифлосланиш хавфини камайтириши мумкинлиги кўрсатилди. [19-21].

5. Бактерияларни аниқлаш усуллари: Қон қўйишдан олдин қон таркибий қисмлардан бактерияларни аниқлашнинг турли усуллари, шу жумладан автоматлаштирилган бактерияларни усули (BacT / ALERT tizimi, bioMérieux), тўғридан-тўғри бактерияларни бўйаш, бактериал эндотоксин ва рибосомал тахлиллар, нуклеин кислоталарни синовдан ўтказиш. Бактериал DNK va O2 истеъмол қилиш ёки CO₂ ишлаб чиқариш ўлчовлари учун (Pall BDS, Pall Corporation) [22-25].

МУҲОКАМА ВА НАТИЖАЛАР

Шу билан бирга, ушбу аниқлаш усулларидан хеч бири барча бактериал ифлосланишларни тўлиқ аниқламайди ва қўшимча бактериал скрининг текширувлари, шунингдек бактериал текширувни яхшироқ ўтказиш вақти (яни қон қўйиш вақтига яқинроқ) бактериялар билан ифлосланган қон махсулотларини тўғри аниқлаш эҳтимолини янада ошириш учун керак бўлиши мумкин.

Патогенларни камайтириш усуллари: Патогенларни камайтириш трансфузион трансмиссив инфекцияни хавфини янада камайтириш учун фаол ёндашув бўлиб, кўпчиликка маълум бўлган ва пайдо бўлаётган патогенлар учун самарали бўлиши мумкин. Патогенни инактиватсия қилишнинг мақсади қон

махсулотининг терапевтик самарадорлигини пасайтирмасдан ёки иккиламчи хавфларни келтириб чиқармайдиган трансмиссив патогенларни (бактериялар, вирус ва протоз) камайтиришдир. Қон таркибий қисмларида патогенни инактивация қилиш тушунчаси маълум патогенларнинг қолдиқ хавфини камайтириш ва янги, аммо номаълум патогенларни самарали йўқ қилишдан иборат. Патогенларни камайтириш ёндашувини танлаш унинг қон қуиши учун компонентларини, масалан, қизил қон таркибий қисмлари, тромбоцитлар концентратлари ва плазма ёки плазмадан ишлаб чиқариладиган махсулотларни даволашда ишлатилишига боғлиқ. Юқори харажатлар, баязи қўшимчаларнинг узоқ муддатли ён таъсири ва спора хосил қилувчи бактериялар каби баязи бир патогенларни заарсизлантирилишига қодир эмаслиги каби мумкин бўлган чекловларни енгиб ўтиш керак, бу патогенларсиз қон махсулотларини кенг миқёсида ва арzon нархларда таъминлаш учун керак бўлади.

ХУЛОСА

Қон таъминотига кирадиган юқумли касалликларнинг таҳиди турғун эмас ва янги патогенлар пайдо бўлиши ёки эскилар эпидемиологик кўринишини ўзгартириши билан ривожланиши мумкин. Шунга қарамай, тобора ортиб бораётган глобал талабларга жавоб берадиган хавфсиз ва арzon қон таъминоти мақсадига трансфузион занжиридаги хар бир босқични мувофиқлаштиришни оптимилаштириш, шу жумладан донорларнинг мувофиқлиги мезонларини диққат билан кўриб чиқиши, ишлов бериш пайтида қатый қоидаларга риоя қилиш орқали эришиш мумкин. Сақлаш, мавжуд скрининг тестларининг мақбул тарзда амалга ошириш, патогеннинг инактивация қилиш усулларидан фойдаланиш ва нихоят хар бир қон қуиши зарурлигини баҳолайдиган эҳтиёткор шифокорларнинг хушёрлиги. Ушбу чора-тадбирлар сезгир ва арzon нархларда аниқлаш ва инактивация қилиш усулларини ишлаб чиқиши ва амалга ошириш билан бир қаторда, қон қуиши хатарларни хозирги кунгача мухим деб хисобланиши керак бўлган жойларда хам хавфсизроқ терапия турига айлантириши мумкин.

REFERENCES

1. Pittman M: A study of bacteria implicated in transfusion reactions and of bacteria isolated from blood products. J Lab Clin Med. 1953, 42 (2): 273-288.
2. McEntegart MG: Dangerous contaminants in stored blood. Lancet. 1956, 271 (6949): 909-911. 10.1016/S0140-6736(56)90378-6.

3. Snyder EL, Dodd RY: Reducing the risk of blood transfusion. *Hematology (Am Soc Hematol Educ Program)*. 2001, 433-442.
4. Andreu G, Morel P, Forestier F, Debeir J, Rebibo D, Janvier G, Herve P: Hemovigilance network in France: organization and analysis of immediate transfusion incident reports from 1994 to 1998. *Transfusion*. 2002, 42 (10): 1356-1364. 10.1046/j.1537-2995.2002.00202.x.
5. Faber JC: Haemovigilance procedure in transfusion medicine. *Hematol J*. 2004, 5 Suppl 3: S74-82. 10.1038/sj.thj.6200427.
6. Faber JC: Worldwide overview of existing haemovigilance systems. *Transfus Apher Sci*. 2004, 31 (2): 99-110. 10.1016/j.transci.2004.07.004.
7. Wagner SJ: Transfusion-transmitted bacterial infection: risks, sources and interventions. *Vox Sang*. 2004, 86 (3): 157-163. 10.1111/j.0042-9007.2004.00410.x.
8. Hillyer CD, Josephson CD, Blajchman MA, Vostal JG, Epstein JS, Goodman JL: Bacterial contamination of blood components: risks, strategies, and regulation: joint ASH and AABB educational session in transfusion medicine. *Hematology (Am Soc Hematol Educ Program)*. 2003, 575-589.
9. CDC: Fatal Bacterial Infections Associated with Platelet Transfusion-United States, 2004. *MMWR Morb Mortal Wkly Rep*. 2005, 54 (7): 168-170.
10. Brecher ME, Hay SN: Bacterial contamination of blood components. *Clin Microbiol Rev*. 2005, 18 (1): 195-204. 10.1128/CMR.18.1.195-204.2005.
11. Blajchman MA, Goldman M, Baeza F: Improving the bacteriological safety of platelet transfusions. *Transfus Med Rev*. 2004, 18 (1): 11-24. 10.1016/j.tmrv.2003.10.002.
12. American Association of Blood Banks: Guidance on implementation of new bacteria reduction and detection standards . *AABB Bulletin*. 2004, 04 (07):
13. Fang CT, Chambers LA, Kennedy J, Strupp A, Fucci MC, Janas JA, Tang Y, Hapip CA, Lawrence TB, Dodd RY: Detection of bacterial contamination in apheresis platelet products: American Red Cross experience, 2004. *Transfusion*. 2005, 45 (12): 1845-1852. 10.1111/j.1537-2995.2005.00650.x.
14. Niu MT, Knippen M, Simmons L, Holness LG: Transfusion-transmitted Klebsiella pneumoniae fatalities, 1995 to 2004. *Transfus Med Rev*. 2006, 20 (2): 149-157. 10.1016/j.tmrv.2005.11.007.
15. Hogman CF, Engstrand L: Serious bacterial complications from blood components--how do they occur?. *Transfus Med*. 1998, 8 (1): 1-3. 10.1046/j.1365-3148.1998.00118.x.

16. Goldman M, Roy G, Frechette N, Decary F, Massicotte L, Delage G: Evaluation of donor skin disinfection methods. *Transfusion.* 1997, 37 (3): 309-312. 10.1046/j.1537-2995.1997.37397240214.x.
17. McDonald CP, Lowe P, Roy A, Robbins S, Hartley S, Harrison JF, Slopecki A, Verlander N, Barbara JA: Evaluation of donor arm disinfection techniques. *Vox Sang.* 2001, 80 (3): 135-141. 10.1046/j.1423-0410.2001.00029.x.
18. Lee CK, Ho PL, Chan NK, Mak A, Hong J, Lin CK: Impact of donor arm skin disinfection on the bacterial contamination rate of platelet concentrates. *Vox Sang.* 2002, 83 (3): 204-208. 10.1046/j.1423-0410.2002.00219.x.
19. Wagner SJ, Robinette D, Friedman LI, Miripol J: Diversion of initial blood flow to prevent whole-blood contamination by skin surface bacteria: an in vitro model. *Transfusion.* 2000, 40 (3): 335-338. 10.1046/j.1537-2995.2000.40030335.x.
20. de Korte D, Marcelis JH, Verhoeven AJ, Soeterboek AM: Diversion of first blood volume results in a reduction of bacterial contamination for whole-blood collections. *Vox Sang.* 2002, 83 (1): 13-16. 10.1046/j.1423-0410.2002.00189.x.
21. Chassaigne M, Vassort-Bruneau C, Allouch P, Audurier A, Boulard G, Grosdhomme F, Noel L, Gulian C, Janus G, Perez P: Reduction of bacterial load by predonation sampling. *Transfus Apher Sci.* 2001, 24 (3): 253-10.1016/S1473-0502(01)00066-0.
22. Brecher ME, Hay SN, Rose AD, Rothenberg SJ: Evaluation of BacT/ALERT plastic culture bottles for use in testing pooled whole blood-derived leukoreduced platelet-rich plasma platelets with a single contaminated unit. *Transfusion.* 2005, 45 (9): 1512-1517. 10.1111/j.1537-2995.2005.00563.x.
23. Brecher ME, Hay SN, Rothenberg SJ: Validation of BacT/ALERT plastic culture bottles for use in testing of whole-blood-derived leukoreduced platelet-rich-plasma-derived platelets. *Transfusion.* 2004, 44 (8): 1174-1178. 10.1111/j.1537-2995.2004.04033.x.
24. McDonald CP, Roy A, Lowe P, Robbins S, Hartley S, Barbara JA: Evaluation of the BacT/Alert automated blood culture system for detecting bacteria and measuring their growth kinetics in leucodepleted and non-leucodepleted platelet concentrates. *Vox Sang.* 2001, 81 (3): 154-160. 10.1046/j.0042-9007.2001.00104.x.
25. Ortolano GA, Freundlich LF, Holme S, Russell RL, Cortus MA, Wilkins K, Nomura H, Chong C, Carmen R, Capetandes A, Wenz B: Detection of bacteria in WBC-reduced PLT concentrates using percent oxygen as a marker for bacteria growth. *Transfusion.* 2003, 43 (9): 1276-1285. 10.1046/j.1537-2995.2003.00487.