

ISSN: 2181-4007

www.tnmu.uz


THE JOURNAL

OF HUMANITIES & NATURAL SCIENCES

GUMANITAR VA TABIIY FANLAR JURNALI

ISSUE 8
VOLUME I

2024



Informing scientific practices around the world through research and development



TIBBIYOT
NASHRIYOTI
MATBAA UYI

**Gumanitar va
tabiiy fanlar
jurnali**



**Journal of
humanities &
natural sciences**

ЖУРНАЛ ГУМАНИТАРНЫХ И ЕСТЕСТВЕННЫХ НАУК

ISSN: 2181-4007 (print)

№ 8 (03), 2024. Vol. 1

Jurnal O'zbekiston Respublikasi Prezidenti Administratsiyasi huzuridagi Axborot va ommaviy kommunikatsiyalar agentligi tomonidan ro'yxatdan o'tkazilgan (guvohnoma № 040226).

Jurnal O'zbekiston Respublikasi Oliy attestatsiya komissiyasi tomonidan 2023 yil 1 iyundan tibbiyot fanlari bo'yicha dissertatsiyalar asosiy ilmiy natijalarini chop etish tavsiya etilgan ilmiy nashrlar ro'yxatiga kiritilgan (2023 yildagi 01-07/710/1-sonli xat).

Журнал зарегистрирован Агентством информации и массовых коммуникаций при Администрации Президента Республики Узбекистан (свидетельство № 040226).

Журнал включен в перечень научных изданий, рекомендованных к публикации основных научных результатов диссертаций по медицинским наукам с 1 июня 2023 года Высшей аттестационной комиссией Республики Узбекистан (письмо № 01-07/710/1 от 2023).

TIBBIYOT NASHRIYOTI MATBAA UYI
O'zbekiston Respublikasi. Toshkent shaxri. Olmazor tumani. Farobiy ko'chasi - 2. 100109
Tel.: (+998-91) 164-24-40, (+998-71) 214-90-164,
vebsayt: www.tmmu.uz, e-mail: asmehrid@gmail.com

TAHRIRIYAT JAMOASI

BOSH MUHARRIR:

D.Sc., professor
Gaybullayev Asilbek Asadovich

TAHRIRIYAT RAISI:

D.Sc., professor
Madazimov Madamin Muminovich

BOSH MUHARRIR O'RINBOSARI

D.Sc., professor
Teshayev Oktyabr Ruxillaevich

MA'SUL KOTIB

Niyazova Zebiniso Anvarovna, Ph.D
Xegay Lyubov Nikolaevna, t.f.n.

TAHRIRIYAT HAY'ATI A'ZOLARI

D.Sc., professor Pavalkis Dainius (Litva)	t.f.f.d., dotsent Iskandarov Sherzod Abdig'anievich (O'zbekiston)
D.Sc., professor Megalhayz Tereza (Portugaliya)	t.f.n., dotsent Kadomseva Larisa Viktorovna (O'zbekiston)
D.Sc., professor Syed Naqi Abbas (Hindiston)	Ph.D., dotsent Karimdjanova Guzal Akmal'djanovna (O'zbekiston)
D.Sc., professor Ayji Mano (Yaponiya)	Ph.D., professor Akramova Nozima Akramovna (O'zbekiston)
D.Sc., professor Boymurodov Shuhrat Abdujalilovich (O'zbekiston)	Ph.D., professor Gaybullayev Elbek Azizbekovich (O'zbekiston)
D.Sc., professor Gavrilova Tatyana Valeryevna (Rossiya)	Ph.D., dotsent Iriskulova Elmira Uraimkulovna (O'zbekiston)
D.Sc., professor Ergashev Ulug'bek Yusufjonovich (O'zbekiston)	Ph.D., dotsent Abdullayeva Shakhlo Kurbanburiyevna (O'zbekiston)
D.Sc., professor Ruziev Sherzod Ibodullaevich (O'zbekiston)	f-m.f.n., dotsent Bazarbaev Muratali Irisalievich (O'zbekiston)
D.Sc., professor Nazarov Azadbek Axmedovich (O'zbekiston)	f.f.n., dotsent Oltiev Temir Jonimboevich (O'zbekiston)
D.Sc., professor Muftaydinov Kiyomidin Xamdamovich (O'zbekiston)	Ph.D., dotsent Ergashev Jamol Djuraboyevich (O'zbekiston)
D.Sc., professor Nikonova Lyudmila Ivanovna (Rossiya)	Ph.D., dotsent Xalillaev Adilbek Kurambaevich
D.Sc., dotsent Zufarov Aziz Alimjanovich (O'zbekiston)	Ph.D. Nuraliev Farid Nekkadamovich
D.Sc., dotsent Salaxiddinov Kamoliddin Zuxriddinovich (O'zbekiston)	t.f.n. Muftaydinova Shaxnoza Kiyomiddinovna (O'zbekiston)
D.Sc., dotsent Baxronova Dilrabo Keldiyorovna (O'zbekiston)	Ph.D. Turamuratova Iroda Ilxombaevna (O'zbekiston)
D.Sc., dotsent Usmanova Durdona Djurabaevna (O'zbekiston)	Ph.D. Ismailova Jadida Axmedjanovna (O'zbekiston)
D.Sc., dotsent Omonova Umida Tulkinovna (O'zbekiston)	Ph.D. Abidov Xasanxodja Alisherovich (O'zbekiston)
D.Sc., dotsent Daminova Kamola Maratovna (O'zbekiston)	Ph.D. Jo'raev Abdunazar Xatamnazarovich (O'zbekiston)
D.Sc., dotsent Velázquez Virna Vilchis (Meksika)	Ph.D. Babaraximova Sayyora Boriyevna (O'zbekiston)
D.Sc., dotsent José Ramón González García (Meksika)	Ph.D. Akromov Ulug'bek Sharobiddinovich (O'zbekiston)
D.Sc. Mahkamova Dilbar Kamaldjanovna	Ph.D. Uzbekov Timur Sakenovich (O'zbekiston)
t.f.n., dotsent Daminova Malika Nasirovna (O'zbekiston)	Ph.D. Ismailova Mahfuza Ubaydullaevna (O'zbekiston)

МУНДАРИЖА – ОГЛАВЛЕНИЕ – CONTENTS

МЕДИЦИНА И ЗДРАВООХРАНЕНИЕ

Матмуротов К.Ж., Кучкаров А.А. / Гнойно-воспалительные заболевания брюшинного пространства: разновидность и способы хирургического лечения	6
Ernazarov O.G., Boymuradov Sh.A., Mirzaev A.U. / Uch shoxli nerv nevrалgiyasining etiologiyasi va patogenezi	13
Исламова Н.Б. / Пародонтит касаллигида организмдаги умумий ва маҳаллий ўзгарган кўрсаткичларнинг таҳлили	23
Асранкулова Д.Б., Мамасолиев Н.С., Мамадалиева М.М. / Ҳомиладорларда темир танқислиги профилактика йуллари ва туғруқдан кейинги ёрдам кўрсатиш асослари	28
Мамajanova Sh.K. / Studying the effectiveness of magnesium in the treatment of algodysmenorrhea	31
Muminov B.A., Matmurodov R.J., Xalimova X.M., Umirova S.M. / COVID-19 o'tkazgan bemorlarda parkinson kasalligi motor buzilishlarining interleykin-6 va ferritin miqdori bilan solishtirma tahlili	33
Yusupova U.M. / Modern approach to the treatment of the causes of pregnancy complications in women with ovarian apoplexy	37
Bozorova M.I., Rasulov A.A., Ashurov D.M. / Ni (II) ning norflokasin bilan yangi kompleks birikmasi sintezi va uning analiz natijalari tahlili	41
Bazarbayev M.I., Qiyomova M.M. / Ultra keng polosali optik kogerent elastografiya yordamida olingan materiallar va to'qimalarning mexanik xususiyatlarini tahlil qilish algoritmini neyron tarmoqlari yordamida optimallashtirish.....	47
Асранкулова Д.Б., Мамасолиев Н.С., Абдукахорова С.К., Мамасолиев З.Н. / Изучение проблем, связанных с физической мультиморбидностью (патогенетические, клинические и эпидемиологические аспекты у беременных).....	56
Ҳамдамов Б.З., Рахимов А.Ш. / Диабетик тоvon синдромида йирингли-некротик асоратлар патогенези	60
Тешаев О.Р., Исломов А.А. / Диабетик оёқ синдромининг йирингли-некротик кўринишларини даволашда жарроҳлик ёндашувларни такомиллаштириш	65
Наврүзова Ш.И., Артикова М.М. / Клинические особенности и коморбидность при хронических вирусных гепатитах.....	70
Ахматова Г.Р., Наврүзова Ш.И. / Прогнозирование цервикальных заболеваний, ассоциированных вирусом папилломы человека.....	75
Ахмедова Н.М. / Клиническая характеристика наружного генитального эндометриоза	79
Хайдарова Д.К., Сафаров К.К. / Современные представления о клинической картине и когнитивных нарушениях при болезни паркинсона.....	83
Ихтиярова Г.А., Розикова Д.К. / Влияние ретрохориальной гематомы на исход беременности у пациентов с угрожающим выкидышем	89
Тешаев Ш.Ж., Эшонқулов Ж.О. / Аҳолини маиший ва ичимлик таъминоти учун фойдаланиладиган сув манбалари. Сув таркибининг инсон организмга таъсирини таҳлил қилиш	92
Тешаев Ш.Ж., Озодов Ж.Х. / Татуировка билан боғлиқ муаммолар ва замонавий ечими усулларини ўрганиш.....	97

Султанова Л.Д., Шеров Ж.А. / Ўткир нурланиш таъсири остида оқ зотсиз каламушларда юрак морфологик ўзгаришлари	103
Султанова Л.Д., Кежаев З.М. / Ўткир нурланиш таъсирида юракдаги морфо-функционал ўзгаришларни даво-профилактикасига янгича ёндашув	108
Усмонов И.Х., Искандарова И.М. / Хавфли ўсмалар ва ўпка туберкулёзини даволашда клиник ва диагностик хусусиятлар ҳамда қийинчиликлар	113
Эшбаев Э.А., Исақов Қ.Б. / Цереброваскуляр касалликларда бош мия ичи қон томирларининг гистокимёвий ўзгаришлари	118
Султанова Л.Д., Адизов И.Ш. / Болаларда Неходжкин лимфомаларнинг клиник-иммунологик хусусиятлари ҳамда даволашни оптималлаштиришга қаратилган замонавий қарашлар	123
Фаттахов Р.А., Хасанова Л.Э., Нарова Н.Э., Юнусходжаева М.К. / Результаты исследования функциональных свойств ротовой жидкости у лиц с зубочелюстными аномалиями	128
Кузиева Г.А. / Проведение предварительной диагностики, профилактики и предупреждения медицинских проблем (при гипертензивных и вегетативных кризах при беременности)	131
Тешаев Ш.Д., Шарипова М.А. / Зотсиз каламушнинг юрак физиологияси ва юрак етишмовчилигининг патофизиологиясини таҳлил қилиш	135
Идиев Г.Э., Бабаева Н.М. / Эффективность применения фотодинамической терапии в комплексном лечении тяжелых форм лейкоплакии слизистой оболочки полости рта	141
Нуров Н.Б. / Кекса ёшдаги беморларда тўлиқ ва қисман адентияга ортопедик стоматологик ёрдам кўрсатилишини такомиллаштириш	145
Хасанова Д.А., Жумаева М.М. / Ультразвуковые и морфологические исследования стенок желчного пузыря при холециститах	151
Uterova G.B., Muxsinova M.X., Azimov Sh.T., Vahabova G.A., Ubaydullayeva O.X. / Bolalarda marfan sindromining klinik hususiyatlari	157
Нуралиева У.М. / Развитие дисбиоза кишечника у тимэктомированных лабораторных животных в эксперименте	160
Гаффоров С.А., Мадаминова Н.С., Астанов О.М. / Особенности оказания ортопедической стоматологической помощи у больных с психиатрическими патологиями	165
Раджабов Ш.Ж., Наврузова Ш.И. / Структура и клинические особенности urgentных состояний при острых респираторных заболеваниях у детей	174
Ражабова Д.Д., Наврузова Ш.И. / Биохимические показатели крови при холестатическом синдроме ..	179
Rasulova S.H., Navruzova Sh.I., Muxamedova Sh.T. / Bolalarda allergiya va ta'sirlangan ichak sindromining differential tashhisi	184
Тешаев Ш.Ж., Барноев Р.И. / Азот диоксидидан келиб чиққан ўпка фиброзининг қизилўнгач деворидаги морфологик ўзгаришларни мувофиқлаштириш	189
Алаудинов С.З., Казакова Н.Н. / Бронхиал астма билан оғриган беморларнинг умумий тавсифи ва фойдаланилган тадқиқот усуллари	197
Ходжиева Д.Т., Гафарова С.С. / I тип қандли диабетдаги диабетик полиневропатия ривожланишида иммун бузилишлар ва генетик омиллар	202

Базарбаев М.И., Элмуротова Д.Б., Нематов Ш.К., Азимов Ш.Ш., Дамиров Т.З., Махкамов А.Р. / Современные подходы к гигиене рук медицинского персонала.....	208
Марасулов А.Ф., Абдуганиева Ш.Х. / Об опыте преподавания математики и информатики в медицинских ВУЗах	218
Галиева З.И., Иноятова Ф.Х., Таджиева Х.С., Милушева Р.Ю., Рахманова В. / Влияние низкомолекулярного хитозана на течение экспериментального острого токсического гепатита	221
Хасанов М.М., Номанов А.А., Мухтаров Ш.Т. / Применение модели прогнозирования осложненного течения суправезикальной обструкции у пациентов с мочекаменной болезнью в клинической практике	232
Бобоев К.Т., Каримов Х.Я., Мусашайхова Ш.М., Мусашайхов Х.Т. / Клинические и молекулярно- генетические особенности формирования острого миелоидного лейкоза	237
Ирисметов М.Э., Хошимов Ж.Т., Таджимуродов М.Б., Режабов Қ.Н. / Орқа бутсимон боғлам жароҳати диагностикаси, патогенези ва замонавий даволаш усули	240
Teshaev Sh.J., Ochilova D.A. / Daun sindromining kelib chiqish sabablariga zamonaviy qarashlar.....	247
Султанова Л.Д., Жумаев Ф.А. / Спастика тетрапарез мавжуд болаларда, болаликнинг I-II давридаги антропометрик текширув усулларига қаратилган замонавий ёндашув.....	252
Хамдамов Б.З., Ташева Ф.О. / Хроническая обструктивная болезнь легких: обзор заболевания в свете современных представлений.....	258
Ходжаматов И.Ю., Мирзамуродов Х.Х. / Современное состояние вопроса полисегментарного остеосинтеза нижних конечностей по илизарову с целью устранения деформаций и укорочений у детей и подростков.....	263
Хамдамова Г.Р., Наврузова Ш.И. / Диагностика аллергической сенсибилизации при дермо- респираторном синдроме	270
Ходжаева Ш.Ш., Ражабов О.А. / Воздействие патологии желудочно-кишечного тракта на формирование воспалительных заболеваний тканей пародонта	275
Хожаназарова С.Ж., Юсуфова М.А. / Пестицидлар фонида туғилган авлод каламуш буйраги постнатал онтогенезининг морфометрик ўзгаришлари	282
Ядгарова Г.С. / Комплексная оценка и улучшение качества жизни детей с врожденными расщелинами губы и неба.....	290
Мираюбова Ш.Р., Хамракулова М.А. / Гигиеническая оценка условий труда работников основных профессий гидротехнических сооружениях Урта-Чирчикского ГЭС.....	293
Марасулов А.Ф. / Формирование умений студентов строить и исследовать количественные модели ..	301
Alimova D.D. / Therapeutic-diagnostic algorithm in different phenotypes of polypous rhinosinusitis	308
Қосимов О.Ш., Бектимиров А.М.-Т., Таджиева Н.У., Абдуллаев А.О. / Экспериментал ҳайвонлардан олинган поливалент диагностик ичак иерсиниози зардобларининг махсуслигини гомологик микробиологларда ўрганиш.....	314

COVID-19 O'TKAZGAN BEMORLARDA PARKINSON KASALLIGI MOTOR BUZILISHLARINING INTERLEYKIN-6 VA FERRITIN MIQDORI BILAN SOLISHTIRMA TAHLILI

Muminov Bekzod Askarovich - tayanch doktorant
Matmurodov Rustambek Jumanazarovich - t.f.d., professor
Xalimova Xanifa Muxsinovna - t.f.d., professor
Toshkent Tibbiyot Akademiyas (Toshkent, O'zbekiston)
Umirova Surayyo Mamurjonovna - Ph.D., assistant
Samarqand Davlat Tibbiyot Universiteti (Samarqand, O'zbekiston)

Rezyume. Maqolada Covid-19 o'tkazgan Parkinson kasalligi bilan og'rikan bemorlarda motor buzilishlarining interleykin-6 va ferritin miqdori bilan solishtirma tahlili natijalari keltirilgan. UPDRS shkalasining motor jihatlari, kundlik harakatlar va motor asoratlar ko'rsatkichlari bo'yicha IL-6 miqdorining va ferritinemiya darajasining oshib borishi teskari korrelyatsion bog'liqlikka ega. IL-6 miqdorining va ferritinemiya darajasining oshib borishi esa o'zaro to'g'ri korrelyatsion bog'liqlikka egadir.

Kalit so'zlar: Covid-19, Parkinson kasalligi, interleykin-6, ferritin.

СРАВНИТЕЛЬНЫЙ АНАЛИЗ МОТОРНЫХ НАРУШЕНИЙ БОЛЕЗНИ ПАРКИНСОНА С УРОВНЯМИ ИНТЕРЛЕЙКИНА-6 И ФЕРРИТИНА У ПАЦИЕНТОВ ПЕРЕНЕСШИХ COVID-19

Муминов Бекзод Аскарлович - базовый докторант
Матмуродов Рустамбек Жуманазарович - д.м.н., профессор
Халимова Ханифа Мухсиновна - д.м.н., профессор
Ташкентская Медицинская Академия (Ташкент, Узбекистан)
Умирова Сурайё Мамуржоновна - Ph.D., ассистент
Самаркандский Государственный Медицинский Университет (Самарканд, Узбекистан)

Резюме. В статье представлены результаты сравнительного анализа уровня интерлейкина-6 и ферритина с двигательными нарушениями болезнью Паркинсона, у больных перенесших Covid-19. Повышение уровня ИЛ-6 и ферритинемии имеет обратную корреляцию с двигательными аспектами повседневной деятельностью и показателями моторного ассортимента UPDRS. Увеличение количества ИЛ-6 и уровня ферритинемии имеет правильную корреляцию.

Ключевые слова: Covid-19, болезнь Паркинсона, интерлейкин-6, ферритин.

Covid-19 infeksiyasi o'tkir respirator infeksiya hisoblanib, betakoronavirus oilasiga mansub bo'lgan SARS-CoV-2 virusi tomonidan chaqiriladi va zooatroponoz hisoblanadi. SARS-CoV-2 qobiqli bir zanjirli (+) RNK saqlovchi virusdir va birinchi marta 2019-yil dekabr oyida aniqlandi. 2020-yil yanvarda epidemiya ko'rinishida tarqaldi, 2020-yil 11-martda esa Butun dunyo sog'liqni saqlash tashkiloti tomonidan pandemiya deb e'lon qilindi. Kasallik kontakt va havo-tomchi yo'li bilan tez tarqalib, 14 kun inkubatsion davriga ega. SARS-CoV-1 virusga o'xshab, organizmdan tashqarida ob'ekt yuzasiga qarab turib, 3 soatdan 4 kungacha o'z hayot faoliyatini saqlaydi [3,6,7,11,13,16].

Parkinson kasalligi (PK) nasliy degenerativ kasallik hisoblanib, dofaminergik, xolinergik, glutamatergik va serotoninergik neurotransmitterlar faoliyatining buzilishi bilan namoyon bo'ladi. Shuning uchun ham ushbu kasallik motor va motor bo'lmagan buzilishlar bilan kuzatiladi. Motor buzilishlarning asosiy klinik ko'rinishlarini titroq, rigidlik va gipokineziya tashkil qilsa, nomotor buzilishlarga kognitiv buzilishlar, uyqu buzilishlari, hid bilishning buzilishi,

depressiya, vegetativ buzilishlar, ruhiy buzilishlar, xavotirlik va qabziyat kiradi [1,2,4,5,9,15,18].

Roberto C. va hammualliflarning ma'lumotlariga ko'ra (2020-yil) Covid-19 infeksiyasi fonidagi diareya PK uchun maxsus bo'lgan UPDRS II va IV qismlari shkalasini yomonlashtiradi va bu holat kasallik fonidagi diareya rivojlanishi bilan bog'liq. Shuningdek siydik ajralishining buzilishi va holsizlik PKda motor fluktuatsiya rivojlanishiga turtki bo'lishi mumkin. Olimlar Covid-19 infeksiyasi davrida PKning motor va nomotor buzilishlarini og'irlashganini ilgari surdilar.

Covid-19 infeksiyasidan keyingi Parkinson kasalligining klinik kechishini o'rganishga harakat qildilar. Maxsus shkalalar yordamida PKning simptomlarini baholab solishtirib ko'rildi, laborator ma'lumotlarni (interleykin-2, interleykin-13, Ig-G) klinik simptomlar bilan taqqosladilar. Olingan natijalari va xulosalariga ko'ra Covid-19 infeksiyasi neurodegenerativ jarayonni kuchaytirib yuborishi mumkin. Ta'kidlanishicha, Covid-19 infeksiyasini o'tkazgan hamma Parkinson kasalligi bilan og'rikan bemorlarda ham kasallik belgilari bir xilda og'irlash-

maydi, bu organizmning immunologik reaksiyasiga uzviy bog'liqdir [8,10,12,14,17].

Yuqoridaig ma'lumotlarni inobatga olgan holda, biz Covid-19 o'tkazgan bemorlarda parkinson kasalligi motor buzilishlarining interleykin-6 va ferritin miqdori bilan solishtirma tahlil qilishni maqsad qilib oldik.

Tadqiqot materiali va usullari. Parkinson kasalligida interleykin-6 va ferritin miqdorining o'zgarish darajalarini solishtirish uchun umumiy 90 nafar PK bilan og'rigan bemorlarning qon zardobi tekshiruvdan o'tkazildi. PK bilan og'rigan bemorlarning 48 nafari (53,3%) erkak va 42 nafari (46,7%) ayollarni tashkil qildi. PK bilan og'rigan erkak bemorlarning yoshi 22-70 yoshgacha bo'lib, o'rtacha 46,6±9,8 yosh va ayollarning yoshi 28-78 yoshgacha bo'lib, o'rtacha 53,7±11,9 yoshni tashkil etdi. Kasallik davomiyligi erkaklarda 5,1±3,6 yilni tashkil etgan bo'lsa, ayollarda esa 6,5±4,8 yilni tashkil etdi. Bemorlar 2 ta guruhga ajratildi: 1-guruh Covid-19 o'tkazgan PK bilan og'rigan bemorlar 45 nafar (50%) va Covid-19 o'tkazmagan PK bilan og'rigan bemorlar 45 nafar (50%). Bizning tadqiqotimizda qon zardobidagi glial IL-6 va ferritin miqdori immunoferment analiz (IFA) usuli bilan tekshirildi. Buning uchun belgilab olingan bemorlarning bilak vena qon tomiridan 10,0 ml miqdorda qon olindi va 10 min davomida minutiga 2000 marta tezlikda sentrifuga qilindi. Qon zardobi maxsus probirkalarga solinib muzlatgichda muzlatilgan holatda saqlandi. Barcha bemorlarda PR harakatga bog'liq bo'lgan buzilishlar UPDRS shakalsi II, III va IV punktlari bo'yicha solishtirildi. Olingan natijalar statistik tahlil qilindi.

Tadqiqot natijalari. Biz Covid-19 o'tkazgan va o'tkazmagan bemorlarda motor simptomlarning uchrash chastotasini bemorlar qon zardobi tarkibidagi interleykin-6 va ferritinemiya darajasi bilan solishtirma tahlil qildik. Olingan natijalar shuni ko'rsatmoqdaki, 1-guruhdagi koronavirus o'tkazgan bemorlarda UPDRS shkalasi 38,9±4,8 ball va 2-guruhdagi koronavirus o'tkazmagan bemorlarda esa 24,4±2,19 balldan iborat bo'lgan bo'lsa, shunga mos ravishda 1-guruhdagi Covid-19 o'tkazgan bemorlar qon zardobi tarkibida IL-6 miqdori 21,37±5,5 pg/ml ekanligi ma'lum bo'ldi va nazorat guruhga nisbatan ishonchli farq qildi, p≤0,001. 2-guruhdagi Covid-19 o'tkazmagan bemorlar qon zardobi tarkibida IL-6 miqdori esa 12,91±4,7 pg/ml ekanligi ma'lum bo'ldi va nazorat guruhga nisbatan ishonchli farq qildi, p≤0,001. Shuningdeku, 1-guruh, ya'ni Covid-19 o'tkazgan PK bilan og'rigan bemorlar qon zardobi tarkibidagi ferritinemiya 317,42±14,48 ng/ml va 2-guruh, ya'ni Covid-19 o'tkazmagan PK bilan og'rigan bemorlar qon zardobi tarkibida ferritin miqdori 122,4±11,8 ng/ml darajada namoyon bo'ldi, ya'ni UPDRS shkalasi bemorlar zardobi tarkibidagi interleykin-6 miqdori va ferritinemiya darajasi bilan to'g'ri korrelyatsion bog'liqlikka ega bo'ldi. Statistik Cheddok shkalasi bo'yicha solishtirilganda, UPDRS shkalasi interleykin-6 miqdori bilan yuqori manfiy korrelyatsion bog'liqlikka ega ekan (r-0,7-0,9 va natija -0,8, ferritinemiya darajasi bilan esa sezilarli darajada manfiy korrelyatsion bog'liqlikka ega ekan, (r-0,5-0,7 va natija -0,6). Interleykin-6 miqdori va ferritinemiya darajasi esa yuqori darajadagi musbat korrelyatsion bog'liqlikka ega ekan, (r - 0,9-0,99, natija +0,95), 1-jadval.

1-jadval.

UPDRS shkalasi II punkti, interleykin-6 va ferritinemiya darajasining korrelyatsion tahlili.

Guruhlar	UPDRS	IL-6	Ferritinemiya	UPDRS va IL-6	UPDRS va ferritinemiya	IL-6 va ferritinemiya
Covid+	38,9±4,8	21,37±5,5	317,42±14,48	r-0,8	r-0,6	r+0,95
Covid-	24,4±2,1	12,91±4,7	122,4±11,8	r-1,8	r-1,3	r+1,6
Farq	p≤0.001	p≤0.001	p≤0.001			

Tadqiqotning keyingi bosqichida UPDRS shkalasi natijalarini kundalik harakat buzilishlari bo'yicha bemorlar qon zardobi tarkibidagi interleykin miqdori va ferritinemiya darajasi bilan solish-

tirma tahlil qildik. UPDRS III qism - kundalik hayotning harakat jihatlari 16 ta punkt bo'yicha solishtirilganda, ishonchli farq kuzatildi korrelyatsion bog'liqlik aniqlandi, 2-jadval.

2-jadval.

UPDRS shkalasi III punkti, interleykin-6 va ferritinemiya darajasining korrelyatsion tahlili.

Guruhlar	UPDRS	IL-6	Ferritinemiya	UPDRS va IL-6	UPDRS va ferritinemiya	IL-6 va ferritinemiya
Covid+	51,4±11,6	21,37±5,5	317,42±14,48	r-0,7	r-0,66	r+0,9
Covid-	33,7±10,3	12,91±4,7	122,4±11,8	r-1,5	r-1,4	r+1,5
Farq	p≤0.001	p≤0.001	p≤0.001			

Keyingi bosqichda UPDRS shkalasining 4-punkti bo'yicha 6 ta ko'rsatkichlar immunologik tekshirish natijalari bilan solishtirildi. 1-guruhdagi

bemorlarda bemorlarda 18,6±6,6, 2-guruhdagi bemorlarda ushbu ko'rsatkich 7,7±5,3 ballga ega bo'lgan bo'lsa, shunga mos ravishda interleykin-6

miqdori va ferritinemiya daraajasi keskin farqlandi. Motor asoratlar solishtirildi va og'ir daradadagi diskineziyalarning davomiyligi bilan bog'liq bo'lgan buzilishlar, o'rta darajadagi diskineziyalarning funksional ta'siri bilan bog'liq buzilishlar, o'rta va og'ir darajadagi flyuktuatsiya, «o'chirish» bilan bog'liq muammolar, o'rta va og'ir darajadagi flyuktuatsiya funktsi-

onal ta'siri bilan bog'liq muammoli buzilishlar, og'ir darajadagi murakkab motor flyuktuatsiyalar muammolari, o'rta va og'ir darajadagi og'riqli distoniya bilan bog'liq bo'lgan muammolar ko'rinishdagi buzilishlar interlekyin-6 miqdorining oshishi va ferritinemiya darajasining yuqoriligi bilan namoyon bo'ldi, 3-jadval.


3-jadval.

UPDRS shkalasi IV punkti, interlekyin-6 va ferritinemiya darajasining korrelyasion tahlili.

Guruhlar	UPDRS	IL-6	Ferritinemiya	UPDRS va IL-6	UPDRS va ferritinemiya	IL-6 va ferritinemiya
Covid+	18,6±6,6	21,37±5,5	317,42±14,48	r-0,7	r-0,66	r+0,9
Covid-	7,7±2,19	12,91±4,7	122,4±11,8	r-1,5	r-1,4	r+1,5
Farq	p≤0.001	p≤0.001	p≤0.001			

Olingan natijalar bo'yicha shunday xulosaga kelish mumkin, bemorlarda Parkinson kasalligi motor buzilishlarining darajalari bemorlar qon zarfobi tarkibidagi interlekyin-6 va ferritin darajasi bilan uzviy bog'liqdir. UPDRS II, III, IV punktlari

ko'rsatkichlari interlekyin-6 va ferritin darajasi bilan teskari korrelyasion bog'liqlikka ega, interlekyin-6 va ferritin darajasi esa to'g'ri korrelyasion bog'liq bilan namoyon bo'ladi, 1-rasm.



1-rasm. UPDRS II, III, IV punktlari ko'rsatkichlari interlekyin-6 va ferritin darajasi bilan solishtirma tahlili.

Xulosalar. Covid-19 o'tkazgan PK bilan og'ir-gan bemorlarda interlekyin-6 «sitokinli bo'ron»-sababchisi bo'lib, ferritinemiya darajasini oshirib yuboradi, natijada bosh miyaning qora modda tarkibidagi neyrodeneratsiya jarayoni kuchayadi va kasallikning motor belgilari jadallashadi. Kasallikning klinik shakllarining transformatsiyaga uchrashi va quyi bosqichlardan yuqoriga o'tib ketishi immunologik ko'rsatkichlar bilan bog'liqlikka egadir. UPDRS shkalasining motor jihatlari bo'yicha, kundalik harakat bo'yicha va motor asortalar ko'rsatkichlari bo'yicha IL-6 miqdorining va ferritinemiya darajasining oshib borishi teskari korrelyatsion bog'liqlikka ega bo'lsa, IL-6 miqdorining va ferritinemiya darajasining oshib borishi to'g'ri korrelyatsion bog'liqlikka egadir.

ADABIYOTLAR

1. Е.А. Катунина, А.Ю. Лебедева, З.Р. Бекова, А.Р. Погорова. Особенности течения COVID-19 у больных с болезнью Паркинсона (результаты собственных наблюдений). Бюллетень Национального общества по изучению болезни Паркинсона и расстройств движений. DOI: 10.24412/2226-079X-2022-12442, 2022. - №2.- С. 87-90.

2. Залялова З.А., Богданов Э.И., Латыпова Г.Р., Яковлева Л.А. Клинико-эпидемиологические аспекты работы Казанского центра экстрапирамидной патологии // Бюллетень сибирской медицины.- Томск, 2008. - №5. - С. 151-157.

3. Залялова З.А., Хасанова Д.М. Риск развития и течение COVID-19 у пациентов с болезнью Паркинсона. Журн. неврол. и психиатр. им. С.С. Корсакова. 2021;121 (9). - С. 152-156.
4. Захаров В.В. Деменция при болезни Паркинсона // Неврол.Журн. – Москва, 2006. – №1. – С. 13-18.
5. Захаров В.В. Дофаминергическая и норадренергическая терапия когнитивных нарушений // Журнал неврологии и психиатрии. – Москва, 2006. – №9. – С. 34-38.
6. Умирова С.М., Матмуродов Р.Ж., Каттаёшли беморларда диабетик полинейропатияларни эрта ташхислаш ва даволаш хусусиятлари // Тиббиётда янги кун журнали, Бухоро, 6 (44) 2022.- С.162-166.
7. Халимова Х.М., Матмуродов Р.Ж., Умирова С.М., Covid-19 о'тказган беморларда диабетик полинейропатияларнинг динамикасини баҳолаш // Биомедицина ва амалиёт журнали, Tadqiqot.uz., Тошкент-2023, - №3. - С. 310-314.
8. Alipoor SD, Mortaz E, Varahram M, Garsen J, Adcock IM. The immunopathogenesis of neuroinvasive lesions of SARS-CoV-2 infection in COVID-19 patients. *Frontiers in Neurology* 2021 Jul;12. - P. 697079.
9. Artusi CA et al. COVID-19 in Parkinson's disease: report on prevalence and outcome. *Parkinsonism Relat. Disord.* 2020;80. – P. 7-9.
10. El-Qushayri AE et al. The impact of Parkinson's disease on manifestations and outcomes of Covid-19 patients: a systematic review and meta-analysis. *Rev. Med. Virol.* 2022;32(2). – P. 2278.
11. Jin J.M. et al. Gender differences in patients with COVID-19: focus on severity and mortality. *Front. Public Health.* 2020;8. – P. 152.
12. Li Z., Yi Y., Luo X. et al. Development and clinical application of a rapid IgM-IgG combined antibody test for SARS-CoV2 infection diagnosis. *J. Med. Virol.* 2020; 92: 1518–24. doi: 10.1002/jmv.25727.
13. Papa S.M., Brundin P., Fung V.S.C. et al. Impact of the COVID-19 Pandemic on Parkinson's Disease and Movement Disorders. *Movement Disorders* [published online ahead of print, 2020 Apr 6]. PMID: 32250460. DOI: 10.1002/mds.28067.
14. Qin C., Zhou L., Hu Z. et al. Dysregulation of immune response in patients with COVID-19 in Wuhan. China. *Clin Infect Dis.* 2020; 71(15): 762–8. doi: 10.1093/cid/ciaa248.pii: ciaa248.
15. Shalash A et al. Mental health, physical activity, and quality of life in Parkinson's disease during COVID-19 pandemic. *Mov. Disord.* 2020;35(7). – P. 1097-1099.
16. World Health Organization. WHO Coronavirus (COVID-19) Dashboard. Available from: <https://covid19.who.int/> Accessed 2022 May 30.
17. Zhang B., Zhou X., Zhu C. et al. Immune phenotyping based on neutrophil-to-lymphocyte ratio and IgG predicts disease severity and outcome for patients with COVID-19. *medRxiv* [Preprint] 16.03.2020. doi: 10.1101/2020.03.12.20035048.
18. Zheng KS et al. Clinical characteristics of exacerbations in Parkinson disease. *Neurologist.* 2012;18(3):120-4. 5. Brugger F et al. Why is there motor deterioration in Parkinson's disease during systemic infections – a hypothetical view. *NPJ Parkinsons Dis.* 2015;1:15014.

COMPARATIVE ANALYSIS OF MOTOR DISORDERS IN PARKINSON'S DISEASE WITH INTERLEUKIN-6 AND FERRITIN LEVELS IN PATIENTS WHO HAVE RECOVERED COVID-19

Muminov Bekzod Askarovich - Ph.D.-student
Matmurodov Rustambek Jumanazarovich - D.M.Sc., professor
Khalimova Khanifa Mukhsinovna - D.M.Sc., professor
Tashkent Medical Academy (Tashkent, Uzbekistan)
Umirova Surayo Mamurjonovna - Ph.D., assistant
Samarkand State Medical University (Samarkand, Uzbekistan)

Resume. The article presents the results of a comparative analysis of the level of interleukin-6 and ferritin with motor disorders of Parkinson's disease in patients with Covid-19. Increased levels of IL-6 and ferritinemia have an inverse correlation with motor aspects of daily activities and UPDRS motor range scores. The increase in the amount of IL-6 and the level of ferritinemia has a correct correlation.

Key words. Covid-19, Parkinson's disease, interleukin-6, ferritin.

MODERN APPROACH TO THE TREATMENT OF THE CAUSES OF PREGNANCY COMPLICATIONS IN WOMEN WITH OVARIAN APOPLEXY

Yusupova Umida Masharifovna

Andijan State Medical Institute (Tashkent, Uzbekistan)

Resume. Ovarian apoplexy is one of the most common urgent gynecological diseases, diagnosed in 17% -31.7% of patients, the third in the composition of urgent diseases in gynecology, the second among the causes of abdominal bleeding. In the form of AP anemia, which occurs in 40% of patients, abdominal bleeding of varying severity is observed, and urgent surgical treatment is carried out.

This article presents the results of the application of modern methods in the prevention and treatment of complications of ovarian apoplexy.

Key words: apoplexy, ovaries, gynecological diseases, complications.

ТУХУМДОН АПОПЛЕКСИЯСИГА ЧАЛИНГАН АЁЛЛАРДА ҲОМИЛАДОРЛИК АСОРАТЛАРИ КЕЛИБ ЧИҚИШИНING ДАВОЛАШГА ЗАМОНАВИЙ ЁНДАШУВ

Юсупова Умида Машарифовна

Андижон Давлат Тиббиёт Институти (Андижон, Ўзбекистон)

Резюме. Тухумдон апоплексияси кенг тарқалган шошилишч гинекологик касалликлардан бири бўлиб, беморларнинг 17% – 31,7% да ташхис қилинади, гинекологияда шошилишч касалликлар таркибида учинчи, қорин бўшлиғи қон кетишининг сабаблари орасида иккинчи ўринда туради. Беморларнинг 40 фоизида учрайдиган АП нинг анемия шакли билан турли оғирликдаги қорин бўшлиғи қон кетиши кузатилади ва шошилишч жарроҳлик даволаш амалга оширилади.

Ушбу мақолада тухумдон апоплексиясининг асоратларини олдини олиш ва даволашда замонавий усулларни қўллаш натижалари келтирилган.

Калит сўзлар: апоплексия, тухумдон, гинекологик касалликлар, асоратлар.

The relevance of the problem. Despite the fact that the urgent pathology under study cannot be classified as rare, there are a limited number of reports in the modern literature concerning its etiopathogenetic aspects. It is known that the above ovarian pathology is a genetically determined nosological unit, in particular, according to the GPIIIa gene.

According to K. Aalto-Setälä, the presence of the PL-AII allele leads to a high risk of coronary thrombosis, and studies by S. R. Steinhubl, D. J. Moliterno demonstrate a link between the carrier of the PL-AII allele and increased adhesion and aggregation of platelets in arterial vessels. Special attention is paid to the PL-AII allele, the presence of which leads to implantation disorders in the form of insufficient invasion of the trophoblast by surface layers. In the studied ovarian pathology, an increase in the frequency of carrying the PL-AII allele contributes to the violation of the integrity of the ovarian tissue due to weak intercellular contacts in it and increased thrombosis in small ovarian vessels, which is realized in hemorrhage.

In the scientific literature, this nosology is represented by various terms: "ovarian hemorrhage", "rupture of the corpus luteum cyst", "ovarian rupture", "ovarian bleeding" [3]. In gynecological practice, AYA is characterized by sudden hemorrhage

into the ovary and a violation of the integrity of the tissue, and with hemorrhagic form – the development of intra-abdominal bleeding [4-5]. Ovarian apoplexy, according to Russian literature, accounts for 17-19% in the structure of acute gynecological pathology and a third of all emergency hospitalizations in gynecological hospitals [2,6]. Among the causes of peritoneal bleeding, this life-threatening nosology, requiring emergency surgical treatment, ranks second after ectopic pregnancy. AYA relapses are detected in 21.6–69.0% of cases [4].

The majority of scientists agree that sexual intercourse, excessive physical exertion and stress are the main provoking factors of this urgent condition. However, they cannot be the direct cause of ovarian hemorrhage. Acting against the background of hormonal imbalance in the presence of ovarian anatomical changes in the form of a cystic corpus luteum, a multifollicular structure, a persistent follicle and in the presence of sclerosed altered vessels, provoking factors lead to increased pressure in the vessels due to blood filling of the pelvic organs and cause a violation of their integrity.

Hormonal imbalance in the genesis of ovarian apoplexy, according to A. S. Gasparov, A. E. Ter-Ovakimyan, is caused by a stress factor leading to hypercorticism and hyperprolactinemia. Hyperproduc-

tion of these hormones is the cause of the appearance of a multifollicular type of ovarian structure

Among the causes that can lead to a violation of women's reproductive health, gynecological diseases that require surgical treatment play a special role. Ovarian apoplexy as an etiological factor of intra-abdominal bleeding is in second place after ectopic pregnancy, and its prevalence has increased several times in recent decades, despite the existing progress and the use of high-tech methods of diagnosis and treatment. Ovarian apoplexy in 40% of cases requires immediate surgical treatment, leading to resection of part of the organ. In the postoperative period, there is an aggravation of impaired reproductive health, progressive deterioration of hormonal homeostasis, deep circulatory disorder, which are caused not only by ovarian trauma and a decrease in ovarian reserve, but also by the occurrence of adhesions in the pelvic organs.

The main way to restore fertility in patients with PCOS at the moment should be considered assisted reproductive technologies (ART), as a result of which pregnancy is achieved in a specific cycle of ovulation stimulation. The concept of ART includes methods of in vitro fertilization (IVF), as well as conception naturally as a result of various methods of ovulation induction [4, 6].

The successes achieved in the diagnosis and treatment of infertility caused by PCOS have led to an increase in the number of pregnancies and births in this contingent of patients [7].

However, infertility treatment can be considered effective and justified not only at the onset of pregnancy, but with its safe course, with the receipt of healthy, full-fledged offspring, because the priority task of obstetrics is safe motherhood and a "healthy start to life" of future generations.

The study of the features of induced pregnancies is of great practical importance due to the high frequency of complications of the gestational period, perinatal morbidity and mortality.

The purpose of the study. To analyze the scientific literature on the causes and mechanisms of ovarian apoplexy in women living in unfavorable climatic conditions of the subarctic region to prevent the development of relapse of pathology.

Materials and methods of research. To solve the tasks, the study was conducted in 3 stages. At the first stage, a retrospective analysis of 6635 birth histories for 3 years (2021-2023) was carried out in the Andijan AOPC.

Stage II included a laboratory study of the parameters of the hemostasis system of women aged 18 to 35 years (n=82) in the acute period of ovarian apoplexy before surgery: prothrombin time, thrombin time, international normalized ratio, fibrinogen, activated partial thromboplastin time, soluble fibrin-monomer complexes, Willebrand factor, antithrombin III, aggregatogram.

At the III stage, a clinical and laboratory examination of women aged 18 to 35 years (p= P2) who underwent surgical treatment for hemorrhagic ovarian apoplexy was performed. At this stage of the work, the patients were divided into 2 groups (within the framework of a prospective open randomized comparative study in parallel groups).

The results of the study. The results of our study indicated a high incidence of premature amniotic fluid discharge in pregnant women with a history of ovarian apoplexy - 29% vs. 13.3% (p<0.05, OR=2.67) — which is consistent with the literature data. According to modern information sources, the frequency of other complications of childbirth did not exceed the average population indicators. However, in the course of our retrospective analysis, fundamentally different results were obtained. 9.5 times more often, women in labor in the main group had a rapid course of labor - 16.1% versus 1.7% (p<0.05, OR=11.35). In addition, in every third woman, childbirth was complicated by an umbilical cord entanglement (p<0.05, OR=2.9).

According to a number of authors, bleeding in the early postpartum period was observed in isolated cases and did not exceed the average population frequency. According to our data, the blood loss in the postpartum period was significantly greater in the maternity group than in the control group -380 ±40 ml versus 280±20 ml, respectively (p<0.05). In addition, hypotonic bleeding in the early postpartum period was 8.5 times more common in women with a history of ovarian apoplexy - 14.5% versus 1.7% (p<0.05, OR=10.02).

The assessment of newborns on the Apgar scale had no statistically significant differences in the compared groups: in the main group it was 7.61± 0.5 points at the first minute and 7.87± 0.3 points at the fifth minute, in the control group - 7.72± 0.4 and 7.92± 0.5 points, respectively (p>0.05). The neonatal period was characterized by the development of complications such as cerebral ischemia, morpho-functional immaturity, hyperexcitation syndrome, inhibition syndrome, torticollis, cephalomatoma in both groups in equal percentages (p>0.05).

All patients underwent emergency surgical interventions for the anemic form of AYA with the presence of hemoperitoneum from 50 to 600 ml (on average 174.8 +12.9 ml). Laparoscopic signs of chronic salpingitis were found in 28.8% of patients. Adhesive process in the pelvic cavity was diagnosed in 25%. In accordance with the classification of the American Society of Fertility (AFS), 18.3% of patients revealed the I degree of severity of the adhesive process and 6.7% — the III degree.

At the same time, 20% of patients with adhesions had previously undergone surgical interventions on the organs of the abdominal cavity and pelvis, including 6.7% of patients operated in connection with AYA. Comparative analysis of anamnesis data showed no significant differences between pa-

tients of the main and retrospective groups, as well as between patients of subgroups 1A and 1B. Subgroups 1A and 1B did not differ in risk factors for the development of the adhesive process, which made it possible to consider these subgroups adequate for comparison, depending on the use of the enzyme preparation Longidase in complex rehabilitation therapy.

Thus, the analysis of clinical and anamnestic data showed that the vast majority of the examined patients with AYA did not realize reproductive function, and they are interested in preserving it (only 23.8% of patients had childbirth). By the time of surgical intervention, every fourth patient (25%) had already formed an adhesive process in the pelvic cavity (I and III degrees) and almost every third (28.8%) had signs of chronic salpingitis. All patients had factors predisposing to the formation of adhesions, such as the urgency of surgery and the presence of blood in the abdominal cavity.

The above data and literature data on the beginning of the formation of adhesions in the next 36 hours after surgery convince of the validity of the early start of rehabilitation measures aimed at the treatment and prevention of the adhesive process in patients operated for AYA. When choosing the drug Longidase, its prolonged antifibrotic effect, pronounced anti-inflammatory and decongestant effect were taken into account.

The early start of the course from 2 days after surgery pursued the goals: to enhance the effect of anti-inflammatory therapy and to ensure the prevention of adhesions and tuboperitoneal infertility. Morphological examination revealed a corpus luteum cyst with hemorrhage in 91.5% of the patients of the main group, a corpus luteum with hemorrhage in 4.3%, and a rupture of the ovarian follicular cyst in 4.2%.

The obtained results of immunohistochemical examination of ovarian tissue using CD34 antibodies indicate a significant increase in the vascular cross-sectional area (vessel size) of yellow bodies in patients with A (742.33 pixels) compared with those in yellow bodies in patients without A (control group) (320.13 pixels) ($p=0.008$; $r=0.54$). The total area of the vessels in the field of vision in patients with AYA was larger compared to those in the control group due to the larger diameter of the vessels and amounted to 53591.9 pixels versus 23844.2 pixels ($p=0.013$; $r=0.51$). The expression of SAFR and its receptors in the cytoplasm of granulosa-luteal cells and vascular endothelium was assessed as moderate or strong.

The average expression score of SEFR on the evaluation scale in patients with AYA in the vascular endothelium was 1.87, in granulosa-luteal cells — 1.43. In normal yellow bodies, the average expression score was 0.67 and 0.58, respectively. When compared with patients of the control group, we revealed an increase in the expression of the SAFR —

SAFRR1 and SAFRR2 receptors in yellow bodies in patients with AYA. The average expression score of SEFRP1 in patients with AYA was 1.08 in vascular endothelium and 1.62 in granulosa-luteal cells. In patients in the control group, this indicator was 0.42 and 0.75, respectively. The average expression score of SEFRP2 in yellow bodies in patients with AYA was 2.19 in vascular endothelium and 2.08 in granulosa-luteal cells. In the control group, these indicators were 1.42 and 1.25, respectively. Thus, the results obtained showed that during the AYA vascularization of the corpus luteum has a number of features indicating changes in ovarian angiogenesis: the vessels have larger sizes compared to those in the control group, the level of SEFR and its receptors is increased in the ovarian tissue (1 and 2). The average values of the SEFR level in the blood serum taken from patients with A before surgery were 269.2 ± 154.3 pg/ml, in group 1A — 282.6 ± 178.4 pg/ml and in group 1B — 248.1 ± 116.6 pg/ml.

The conducted study indicates the activation of angiogenesis in ovarian tissue in patients with AYA, expressed in a significant increase in the size and area of vessels, as well as in an increase in the expression of SAFR and its receptors both in the endothelium and in granulosa-luteal cells. The revealed features of ovarian angiogenesis complement our ideas about the development of hemorrhage in ovarian tissue in AYA.

The study of systemic angiogenesis after the end of the rehabilitation therapy complex showed that the level of SEFR significantly decreased in the blood serum of patients receiving Longidase for the prevention of adhesions. Since the activation of angiogenesis plays an important role in the formation of adhesions, an increase in the concentration of SAFR in patients who were not prescribed the drug Longidase can be considered as an unfavorable prognostic risk factor for the development of adhesions.

The clinical efficacy of prescribing the drug Longidase as part of complex rehabilitation therapy is confirmed by data on a more favorable course of the postoperative period and rapid normalization of the ultrasound picture, the results of hysterosalpingography, laparoscopy and the frequency of pregnancy compared with those in patients who did not receive the enzyme preparation.

The results obtained convince of the validity of the inclusion in the complex of rehabilitation measures after endosurgical treatment of the drug Longidase, characterized by a prolonged antifibrotic effect, pronounced anti-inflammatory and decongestant action.

Conclusion. Thus, pregnant women who had a history of surgical treatment of ovarian apoplexy, it is advisable to include in the risk group for the development of miscarriage in early terms, complicated gestational course, childbirth and the postpartum period (preeclampsia, gestational anemia, premature discharge of amniotic fluid, entwined umbilical

cord, the development of excessively strong labor with a rapid course of labor, pathological blood loss in childbirth, hypotonic bleeding in the early postpartum period).

REFERENCES

1. Ovarian apoplexy and ruptures of ovarian cysts / A.S. Gasparov et al. - M.: Medical Information Agency, 2018. - 176 p.

2. Wasserlauf, Irina The relative position of chromosomes in the nuclei of ovarian trophocytes in a number of Diptera / Irina Wasserlauf. - M.: LAP Lambert Academic Publishing, 2020. - 300 p.

4. Gasparov, A.S. Ovarian apoplexy and ruptures of ovarian cysts. Monograph / A.S.

3. Gasparov. - M.: Medical Information Agency (MIA), 2022. - 810 p. - 126 p.

4. Diagnosis and treatment of ovarian cancer / Edited by V.A. Gorbunova. - M.: Medical Information Agency, 2020. - 256 p.

5. Kulakov, V. I. Changes in the reproductive system and their correction in women with benign tumors and ovarian tumors / V.I. Kulakov, R.G. Gataulina, G.T. Sukhoi. - M.: Triad-X, 2022. - 256 p.

6. Potin, Vladimir Vsevolodovich Hyperprolactinemic ovarian insufficiency / Potin Vladimir Vsevolodovich. - M.: N-L, 2021. - 904 p.

7. Tsondek, Bernhard Hormones of the ovary and anterior pituitary gland / Bernhard Tsondek. - M.: State Publishing House of Collective farm and State Farm literature, 2022. - 428 p.

СОВРЕМЕННЫЙ ПОДХОД К ЛЕЧЕНИЮ ПРИЧИН ОСЛОЖНЕНИЙ БЕРЕМЕННОСТИ У ЖЕНЩИН С АПОПЛЕКСИЕЙ ЯИЧНИКОВ

Юсупова Умида Машарифовна

Андижанский Государственный Медицинский Институт (Андижан, Узбекистан)

Резюме. Апоплексия яичников - одно из наиболее распространенных неотложных гинекологических заболеваний, диагностируемое у 17% -31,7% пациенток, третье в составе неотложных заболеваний в гинекологии, второе среди причин абдоминальных кровотечений. При форме анемии АП, которая встречается у 40% больных, наблюдается абдоминальное кровотечение различной степени тяжести, и проводится срочное хирургическое лечение.

В данной статье представлены результаты применения современных методов в профилактике и лечении осложнений апоплексии яичников.

Ключевые слова: апоплексия, яичники, гинекологические заболевания, осложнения.