



European journal of science archives conferences series/ Konferenzreihe der europäischen Zeitschrift für Wissenschaftsarchive https://doi.org/10.5281/zenodo.6631546

2 MAY-2022

Institute for Scientific Research and Publication

8

Research and Publishing Center virtualconferences. press

European journal of science archives conferences series/ Konferenzreihe der europäischen Zeitschrift für Wissenschaftsarchive

Internet address: http://ifsrp.edu.ge/

E-mail: info@ifsrp.edu.ge/

Published by Institute for Scientific Research and Publication & Research and Publishing Center

virtualconferences. press

Issued monthly

DOI prefix: https://doi.org/10.5281/zenodo.6631546

Hausarztzentrum Aachen-Forst Trierer Str. 67, 52078 Aachener, Germany

Requirements for the authors.

The manuscript authors must provide reliable results of the work done, as well as an objective judgment on the significance of the study. The data underlying the work should

be presented accurately, without errors. The work should contain enough details and bibliographic references for possible reproduction. False or knowingly erroneous statements

are perceived as unethical behavior and unacceptable.

Authors should make sure that the original work is submitted and, if other authors' works or claims are used, provide appropriate bibliographic references or citations. Plagiarism

can exist in many forms - from representing someone else's work as copyright to copying or paraphrasing significant parts of another's work without attribution, as well as claiming one's rights to the results of another's research. Plagiarism in all forms constitutes unethical

acts and is unacceptable. Responsibility for plagiarism is entirely on the shoulders of the authors. Significant errors in published works. If the author detects significant errors or inaccuracies in the publication, the author must inform the editor of the journal or the publisher about this and interact with them in order to remove the publication as soon as possible or correct errors. If the editor or publisher has received information from a third party that the publication contains significant errors, the author must withdraw the work or correct the errors as soon as possible.

Aachener, Germany 2022

The publisher is not responsible for the materials published in the collection. All materials are Submitted in the author's edition and reflect the personal position of the conference participant.

Contact information of the organizing committee of the conference:

Email: info@ifsrp.edu.ge/ or info@virtualconferences.press

Official site: http://ifsrp.edu.ge/ or www.virtualconferences.press

DOI https://doi.org/10.5281/zenodo.6631546

Available at virtualconferences. press

ORCID 0000-0001-6156-3630

OPEN ACCESS

MEDICINE

Evaluation of antibacterial activeness of aboveground level part of portulaca oleracea l

Fayzullayeva Zamira Rakhmatovna docent, Department of Microbiology

Tashkent Medical Academy Uzbekistan

Khodjaeva Muattar Asadullaevna dotsent, Department of Pharmacognosy

Tashkent Pharmaceutical Institute Uzbekistan

Aim. Scientists of pharmaceutical industry consider herbal plants as a good choice of getting herbal medicines, because, these natural resources struggle against wide range of antibiotic bacteria. Ingredients of medical plants are used in all parts of the world and has characteristics that are used against bacteria, viruses [1]. Utilization of antibiotic was realized from the lowest rate until the highest. In medical science main problem of appearing resistant bacteria comes from incorrect and overuse of antibiotics [2]. Considered urgent resistance of antibiotic has unlimited demand on new and effective therapeutic issues [3]. In this case, there will be need for producing alternative antibacterial medicines using herbals for curing infection related diseases. Portulaca Oleracea L. is considered as a member of Portulacaceae having more than 120 different types. Name Portulaca comes from Latin language "porto" and means "to carry" and "lac" which means milk due to the fact that plant involves milk juice, Oleracea from Latin language meaning "relating to yard" meaning it's use as a vegetable. Plant grows in yards Central Asia and also in all regions of Uzbekistan as useful plant. The use of this plant as a vegetable of medicine was popular from Ancient Egyptian times and in England within the period of Midcentury [4]. It is grown in India East and popular in Europe even nowadays.

Research methods. Dried Portulaca from yard was used as a material of research. 20 gr of plant material was mixed in bottle weighting 200 ml and was added in Soxhlet during 2-5 hours in temperature not increasing above boiling level. Used ingredients: water, methanol, ethanol, at seton, edit and geksan. Ingredients are mixed in 20% dimetil sulfoxide water in order to get ready mixture 100 mg/ml. Ingredients are kept in dark place in 4°C temperature.

53 MAY-2022

Antibacterial activeness was studied in diffusion method. Resistant bacteria were separated from different clinical examples, like, blood, liquid brain. These issues were identified on basis of their morphology, culture and biochemical characteristics and also regarding the test of sensitiveness of antibiotics. These all bacteria were resistant on more than 10 antibiotics. Antibacterial activeness of Portulaca was evaluated with various herbal examples, like, Staphylococcus aureus (ATCC-25923(004134)), Escherichia coli (ATCC-25922(004136)), Pseudomonas aeruginosa (ATCC-27853(004135)), Klebsiella pneumoniae, Proteus mirabilis, Salmonella spp., Enterococcus faecalis, Citrobacter freundii, Acinetobacter baumannii, Streptococcus pneumoniae, Enterococcus faecium and enter. cloacae.

Description of antibacterial activeness of herbal medicine.

Diffusion method was used in order to study antibacterial effectiveness of notcleaned ingredients. During 15 minutes after correcting the mistake in culiat. Sterilised pain was turned into suspensias. Dried upper side of Mueller-Hinton plastin was widened to all bacteria. This is used approximately 6 mm for curing. 20% water dimeticsulfoxid was used as negative control. Procedure is repeated for other microorganisms. Plastin are kept in fridge during 2 hours for full cooling in the temperature of 37₀C during 24 hours. After incubation period diameter of each zone was calculated in millimetres.

Description of minimum ingredients.

In not cleaned ingredients minimum amount of each testing organism was provided in relation with diffusion method. Ingredient (100 mg/ml) in 20% dimetilsulphoxide is added in cleaned water (1:1) in order to reach decreasing level of ingredients of 50 mg/ml until 0.195 mg/ml. Volume of each 100 m kl standardized in culiat (106 KOE/ml). All testing cups combined aerobic on 37₀C temperature during 24 hours and controlled. Minimum using ingredients showed clear zones (8mm) was reached for each testing microorganisms [5, 6].

Research results. Research of antibacterial activeness of ground level Portulaca Oleracea L. showed clear zones of stopping against bacteria (Table №1). The lowest

uses ingredients methanol extract consisted of 0,8 mg/ml used in relation with E. coli, S. aureus and S. pneumonia when providing concentration if 1.5 mg/ml controls S.typhi and E. faecalis (Table №2). Ethanol extract also showed good activeness in relation with bacteria.

Table №1
The antimicrobial activity of Portulaca oleracea L. against multiple drug resistant bacteria

Raw materials	The solvent	Zone of inhibition in mm											
		1	2	3	4	5	6	7	8	9	10	11	12
	Methanol	24	26	18	22	18	22	15	20	18	24	18	16
Leaves	Ethyl	18	20	14	18	14	16	14	18	17	22	16	14
	alcohol												
	Petroleum	12	10	8	8	7	8	9	8	10	10	10	8
	ether												
	Acetone	16	18	12	12	10	12	12	12	11	12	12	11
	n-Hexane	8	8	10	8	9	10	8	8	8	8	8	8

- 1-S. aureus; 2-E. coli; 3-P. aeruginosa; 4-K. pneumoniae; 5-P. mirabilis; 6-S. typhi;
- 7-E. faecalis;
- 8-C. freundii; 9-A. baumannii; 10-S. peumoniae; 11-E. faecium; 12-E. cloacae

Table № 2

Minimum inhibitory concentration of Portulaca oleracea L. against multiple drugresistant bacteria

The solvent	Minimum inhibitory concentration in mg/ml											
Sorvent	1	2	3	4	5	6	7	8	9	10	11	12
Methanol	0.8	0.8	3.2		3.2		3.2	1.5	1.5		1.5	1.5
Ethyl alcohol	1.5	1.5	3.2	6,2	6,2	1.5	1.5	3.2	1.5	1.5	6,2	6,2

Conclusions. 1. Methanol extract from ground level Portulaca showed high activeness against microorganisms Staphylococcus aureus (23mm) and against bacteria and especially Pseudomonas aeruginosa (19 mm). This may be related with triterpenoid and cartinoid, steroid, cteon and triterpenoid. 2. Received results occurred to be similar to received S. Monroe and others. 3. Ethanol extract also showed good activeness in relation with tested bacteria. 4. Extract from ground level Portulaca can be seen as a resource for getting antimicrobial medicines.

- 1. Hassawi D. and Kharma A. Antimicrobial activity of some medicinal plants against Candida albicans. J. Biological Sci, 2016; 6: 109-114.
- 2. Caldwell R, Lindberg D. (Eds.), Understanding Evolution. University of California Museum of Paleontology, 2017.
- 3. WHO Global Strategy for Containment of Antimicrobial Resistance, 2002.
- 4. Okafor Izuchukwu Azuka, Ayalokunrin Mary B. and Orachu Lovina Abu. A review on Portulaca oleracea (Purslane) plant Its nature and biomedical benefits. International Journal of Biomedical Research. 2014; 05(02).
- 5. Aneja KR, Joshi R, Sharma C. Antimicrobial activity of Dalchini (Cinnamomum zeyla nicum bark) extracts on some dental caries pathogens. J Phar Res, 2019; 2(9):1370-2.
- 6. Nkere CK and Iroegbu CU. Antibacterial screening of the root, seed and stem bark extracts of Picralima nitida. Afr J Biotechnol, 2015; 4: 522-526.

56 MAY-2022