


<http://journal.sapub.org/ajmms>

ISSN: 2165-901X

Volume 6 * Number 6 * December 2016



American Journal of Medicine and Medical Sciences



Scientific & Academic Publishing, USA

American Journal of Medicine and Medical Sciences

p-ISSN: 2165-901X
e-ISSN: 2165-9036
<http://journal.sapub.org/ajmms>

Editorial Board

Michael LaPointe	Indiana University Northwest, USA
Wasim Khan	University College London, UK
Lei Zhang	University of California, Los Angeles, USA
Francesco Rotondi	'San Giuseppe Moscati' Hospital, Italy
Marta Palusinska-Szys	Maria Curie Sklodowska University, Poland
Gerardo Carino	Brown University, USA
Gerald Klein	University of the Pacific, USA
Fei Li	University of Illinois, USA
Harunor Rashid	National Centre for Immunisation Research and Surveillance, Australia
Edmund So	Chi Mei Medical Center, Taiwan
Mehmet Can	Gulhane School of Medicine, Turkey
Suleiman Ambali	University of Ilorin, Nigeria
Adel Bondok	Mansoura University, Egypt
Abdelilah Gounni S	Department of Immunology, Canada
Jone Stanley	Texas A&M University, USA
Tufan Mert	Kahramanmaras Sutcu Imam Univ, Turkey

Purpose

American Journal of Medicine and Medical Sciences aims to promote communication among clinical researchers worldwide. This journal publishes original works within the area of clinical medicine. It is academically peer-reviewed and designed for all doctors, researchers, and health workers regardless of their medical specialties, or type of employment.

Range of Topics

The journal features original research articles as well as review articles in all areas of medicine and medical sciences. Subject areas include (but are not limited to) the following fields:

Anaesthesia, Bioengineering, Biomedical Sciences, Dermatology, Epidemiology and Public Health, Health Informatics, History of Medicine, Health Plan and Administration, Medicine, Neurological Sciences, Ophthalmology, Obstetrics, Occupation Safety and Health, Otolaryngology, Pathology, Psychiatry, Radiology, Surgery.

Minimally Invasive Surgery in Restoring Reproductive Function of Female Infertility Caused by Benign Ovarian Structural Changes

Shukurov Farkhad Ishkulovich

Associate Professor, Department of Obstetrics and Gynecology, Tashkent Medical Academy, Uzbekistan

Abstract **The aim of the study:** to assess the effectiveness of advanced minimally invasive laparoscopic surgical interventions in the treatment of women with infertility due to a benign structural changes of the ovaries. **Material and methods:** A survey of 270 patients with infertility who underwent in operation in gynecology department of obstetric complex № 9 in Tashkent in 2014-2016. The average age of patients was 25.8 ± 0.98 years. Primary infertility suffered 160 (59.2%), the secondary - 110 (40.7%) patients. The duration of infertility was from 2 to 9 years. To clarify the causes of infertility all patients were conducted a comprehensive survey that included clinical, laboratory, hormonal, bimanual, microbiological examination, transvaginal ultrasound scan, hysterosalpingography, laparoscopy and hysteroscopy. Endosurgical treatment of patients with infertility was carried out by two methods: laparoscopic and hysteroscopic. **Results:** As a result of investigation of 120 women (44.4%) identified infertility due to follicular ovarian cysts and 120 (44.4%) of infertility associated with impaired ovulation - polycystic ovary syndrome (PCOS). It should be noted that 80 infertile women (30.0%) mentioned a combination of 2 to 3 factors that violate the reproductive function. During laparoscopy the patients with infertility due to a benign structural changes of the ovaries, revealed various pathological changes in the fallopian tubes: 24 (10%) had adhesions in the fimbria tract, 6 (2.5%) was determined by the obstruction of the fallopian tube in the proximal. 3 patients (1.25%) had determined adhesion process 2 - 3 degrees of severity. When hysteroscopic study, 30 women with benign structural changes of the ovaries 26 (10.8%) of women detected intrauterine pathology: chronic endometritis - 11 (4.6%), synechia in the uterus - 5 (2.1%), endometrial hyperplasia - in 4 (1.6%), foreign bodies of the uterus in the form of silk sutures after cesarean section 6 (2.5%). **Conclusions:** The effectiveness of resection of the ovaries in PCOS is increased 1.5 times compared to electrocauterisation. The restoration of women fertility in with benign structural changes of the ovaries amounted to 87.5%, undergoing endosurgical operation in PCOS and cystectomy. A differentiated approach to the choice of method of surgical intervention depending on the kind of benign structural changes of the ovaries. Combined laparoscopy hysteroscopic study allows to identify and eliminate the causes of uterine infertility.

Keywords Benign structural changes of the ovaries, Laparoscopy, Hysteroscopy, Reproductive function

The development of effective methods of overcoming infertility is now the most important public health and social problem. This is due to a high rate of infertility, its adverse impact on the demographic situation in the country, the psychological climate in the family and social adaptation couple [1, 2]. One of the infertility treatment is the use of endosurgical technology, the pregnancy rate after that ranges from 21% to 30% [3, 4]. Diagnostic value of endoscopic methods in identifying the causes of women is fertility problems is extremely high, reached 97.6%. The effectiveness of surgical treatment of women reproductive

age with various gynecological diseases increased significantly after the introduction into clinical practice of new technologies with the use of endoscopic techniques [5, 6]. However, the controversial issues remain of the use of certain technical methods of surgery, different surgical accesses, a new technical means [7, 8]. In connection with the current debate about the treatment of female infertility due to benign structural changes in the ovaries, it is highly relevant to specify the role and place of endoscopic surgical treatment of patients with infertility in the modern world.

1. The Aim of the Study

The aim of the study was the evaluation of the effectiveness of modern minimally invasive endosurgical

* Corresponding author:

farkhadshukurov@mail.ru (Shukurov Farkhad Ishkulovich)

Published online at <http://journal.sapub.org/ajmms>

Copyright © 2016 Scientific & Academic Publishing. All Rights Reserved

surgical interventions in the treatment of female infertility caused by benign ovarian structural changes.

2. Materials and Methods

A survey of 270 patients with infertility who underwent operation in gynecology department of obstetric complex № 9 in Tashkent in 2014-2016. The average age of patients was 25.8 ± 0.98 years. Primary infertility suffered 160 (59.2%), the secondary - 110 (40.7%) patients. The duration of infertility was from 2 to 9 years. To clarify the causes of infertility all patients were conducted a comprehensive survey that included clinical, laboratory, hormonal, bimanual, microbiological examination, transvaginal ultrasound scan, hysterosalpingography, laparoscopy and hysteroscopy. Endosurgical treatment of patients with infertility was carried out by two methods: laparoscopic and hysteroscopic. Laparoscopic surgery was performed by standard methods using equipment of "Karl Storz". Hysteroscopic surgery was performed using an apparatus «Karl Storz» in liquid simultaneously with or separately laparoscopy. Depending on the used operative method, the patients were divided into 2 groups: all 240 patients had the operation on the ovaries by laparoscopy; other 30 patients had by hysteroscopy. In order to assess the effectiveness of various techniques in reproductive surgery there were used the following indicators according to the survey: frequency of recovery of menstrual function and frequency of uterine pregnancy. The obtained facts were processed with the help of statistical software package "Statistic 6.0".

3. Results and Discussion

As a result of investigation of 120 women (44.4%) identified infertility due to follicular ovarian cysts and 120 (44.4%) of infertility associated with impaired ovulation -

polycystic ovary syndrome (PCOS). It should be noted that 80 infertile women (30.0%) mentioned a combination of 2 to 3 factors that violate the reproductive function. During a laparoscopy was discovered characterized by the following signs PCOS: ovaries of a pearl-gray color, smooth, shiny, taut surface with a characteristic vascular pattern or looped vessels, the presence of undercapsular translucent cysts with a diameter of from 0.3 to 1 sm, dense or uniformly thickened shell. Resection of the ovaries with toothed forceps, found small cysts with a diameter of 0.3 - 2 sm with clear or yellowish contents (number of cysts in each ovary was usually equal to from 10 to 15). During a laparoscopy, follicular ovarian cysts, the tumor was smooth - and thin-walled unilocular formation of rounded, containing transparent light yellow liquid, with a diameter of from 50 to 80 mm, located mostly on the side or posterior from the uterus. When follicular cysts were produced in the husking of the capsule of the cyst - cystectomy, but with PCOS commonly employed by 2 options laparoscopic surgery: electrocauterisare (30 patients) and partial resection of the ovaries (90 patients). The laparoscopic electrocauterisation of polycystic ovaries has been performed with using a monopolar needle electrode. Ovarian electrocautery conducted, as for as possible, all the fine-cystic inclusions. Partial resection of the ovaries was performed in the following way: the ovary for free captured pole toothed forceps, and depending on the size of the incision was made around the tongs - the holders of the distance necessary to remove the desired amount of tissue. Excision of tissue is about 0.5 to 1 sm ovarian tissue. During laparoscopy the patients with infertility due to a benign structural changes of the ovaries, revealed various pathological changes in the fallopian tubes: 24 (10%) had adhesions in the fimbria tract, 6 (2.5%) was determined by the obstruction of the fallopian tube in the proximal. 3 patients (1.25%) had determined adhesion process 2 - 3 degrees of severity. (Fig 1)




Figure 1. Concomitant pathology identified during laparoscopy of patients with benign structural changes of the ovaries

surgical interventions in the treatment of female infertility caused by benign ovarian structural changes.

2. Materials and Methods

A survey of 270 patients with infertility who underwent in operation in gynecology department of obstetric complex № 9 in Tashkent in 2014-2016. The average age of patients was 25.8 ± 0.98 years. Primary infertility suffered 160 (59.2%), the secondary - 110 (40.7%) patients. The duration of infertility was from 2 to 9 years. To clarify the causes of infertility all patients were conducted a comprehensive survey that included clinical, laboratory, hormonal, bimanual, microbiological examination, transvaginal ultrasound scan, hysterosalpingography, laparoscopy and hysteroscopy. Endosurgical treatment of patients with infertility was carried out by two methods: laparoscopic and hysteroscopic. Laparoscopic surgery was performed by standard methods using equipment of "Karl Storz". Hysteroscopic surgery was performed using an apparatus «Karl Storz» in liquid simultaneously with or separately laparoscopy. Depending on the used operative method, the patients were divided into 2 groups: all 240 patients had the operation on the ovaries by laparoscopy; other 30 patients had by hysteroscopy. In order to assess the effectiveness of various techniques in reproductive surgery there were used the following indicators according to the survey: frequency of recovery of menstrual function and frequency of uterine pregnancy. The obtained facts were processed with the help of statistical software package "Statistic 6.0".

3. Results and Discussion

As a result of investigation of 120 women (44.4%) identified infertility due to follicular ovarian cysts and 120 (44.4%) of infertility associated with impaired ovulation -

polycystic ovary syndrome (PCOS). It should be noted that 80 infertile women (30.0%) mentioned a combination of 2 to 3 factors that violate the reproductive function. During a laparoscopy was discovered characterized by the following signs PCOS: ovaries of a pearl-gray color, smooth, shiny, taut surface with a characteristic vascular pattern or looped vessels, the presence of undercapsular translucent cysts with a diameter of from 0.3 to 1 sm, dense or uniformly thickened shell. Resection of the ovaries with toothed forceps, found small cysts with a diameter of 0.3 - 2 sm with clear or yellowish contents (number of cysts in each ovary was usually equal to from 10 to 15). During a laparoscopy, follicular ovarian cysts, the tumor was smooth - and thin-walled unilocular formation of rounded, containing transparent light yellow liquid, with a diameter of from 5 to 80 mm, located mostly on the side or posterior from the uterus. When follicular cysts were produced in the husking of the capsule of the cyst - cystectomy, but with PCOS commonly employed by 2 options laparoscopic surgery: electrocauterisare (30 patients) and partial resection of the ovaries (90 patients). The laparoscopic electrocauterisation of polycystic ovaries has been performed with using a monopolar needle electrode. Ovarian electrocautery conducted, as for as possible, all the fine-cystic inclusions. Partial resection of the ovaries was performed in the following way: the ovary for free captured pole toothed forceps, and depending on the size of the incision was made around the tongs - the holders of the distance necessary to remove the desired amount of tissue. Excision of tissue is about 0.5 to 1 sm ovarian tissue. During laparoscopy the patients with infertility due to a benign structural changes of the ovaries, revealed various pathological changes in the fallopian tubes: 24 (10%) had adhesions in the fimbria tract, 6 (2.5%) was determined by the obstruction of the fallopian tube in the proximal. 3 patients (1.25%) had determined adhesion process 2 - 3 degrees of severity. (Fig 1)




Figure 1. Concomitant pathology identified during laparoscopy of patients with benign structural changes of the ovaries

When hysteroscopic study, 30 women with benign structural changes of the ovaries 26 (10.8%) of women detected intrauterine pathology: chronic endometritis - 11 (4.6%), synechia in the uterus - 5 (2.1%), endometrial hyperplasia - in 4 (1.6%), foreign bodies of the uterus in the form of silk sutures after cesarean section 6 (2.5%) (Figure 2).

Conducting laparoscopic surgeries has reduced the duration of the surgery, reduce bleeding, significantly reduce postoperative pain and duration of hospitalization. Eliminating proximal obstruction of the fallopian tubes was performed by hysteroscopic catheterization of the fallopian tubes. Beside that there was the correction of the concomitant gynecological pathology: restoration of the anatomical relationships of the pelvic organs (lysis of adhesions). With the purpose of preserving ovarian reserve surgery for ovarian pathology was organ-: electrocauterisare ovary, partial resection of the ovaries and husking of ovarian cysts (cystectomy). At presence at patients of concomitant intrauterine pathology hysteroscopic conducted its correction: removal of foreign bodies from the uterine cavity in the form of silk sutures after cesarean section) – in 6

(2.5%), the dissection of adhesions in the uterine cavity – in 5 (2.1%). With proper operation the bleeding from the ovary was minimal. The postoperative period was uneventful. Discharge from hospital was made on the 2nd day. Long-term results of endosurgical operations assessed by conducting a prospective study of women: considered the recovery of menstrual function and the occurrence of intrauterine pregnancy. Uterine pregnancy occurred in a total of 210 women (87.5%). The recovery of menstrual function is assessed according to the presence of biphasic rectal temperature, folliculometry with the help of ultrasound and the recovery of fertility of patients suffering from infertility. In 79.6% of patients in the first days after surgery was noted like a menstrual reaction, which is regarded as a positive response of the endometrium in endocrine adjustment caused by surgery on the ovaries. In addition to, we evaluated the results of surgical treatment for 120 patients with PCOS, depending on the type of surgical intervention (Figure 3). The total number of occurrence of intrauterine pregnancy after partial resection of the ovaries was 1.5 times higher than after electrocauterisation of the ovaries amounted to 56.3% and 36.3%, respectively.




Figure 2. The patients with benign structural changes of the ovaries were identified types of pathology during hysteroscopy




Figure 3. The pregnancy rate in patients with PCOS, depending on the type of surgery on the ovaries

Laparoscopy is more efficient regarding the treatment of benign structural changes of the ovaries. This is due to the safety of minimally invasive endoscopic interventions, with minimal drug load in the form of medicines and transfusions, less duration of hospital stay, faster recovery, cosmetic effect, satisfaction of the patient. According to our data, the use of hysteroscopic intervention in conjunction with laparoscopy facilitates a complete diagnosis of the causes of infertility, their simultaneous correction using minimally invasive techniques, which increases the frequency of in womens pregnancy.

4. Conclusions

1. To increase the effectiveness of treatment of patients with infertility due to a benign structural changes of the ovaries, early detection of diseases and the use of minimally invasive surgical procedures in the early stages of the disease.
2. The effectiveness of resection of the ovaries in PCOS is increased 1.5 times compared to electrocauterisation. The restoration of women fertility in with benign structural changes of the ovaries amounted to 87.5%, undergoing endosurgical operation in PCOS and cystectomy.
3. A differentiated approach to the choice of method of surgical intervention depending on the kind of benign structural changes of the ovaries. Combined laparoscopy hysteroscopic study allows to identify and eliminate the causes of uterine infertility.
4. Hysteroscopy is a highly effective method for diagnosing intrauterine womens pathology of reproductive age and can be used as a screening method of examination in women with infertility caused by structural changes of the ovaries, in some cases, as surgery without additional traumatization of the cervical canal and cervix.

REFERENCES

- [1] Adamyan L. V. Minimally invasive surgery in gynecologic practice, gynecology and obstetrics at the. and gynecol. - 2006, app. - S. 11-17.
- [2] Immunohistochemical techniques: a Manual / Ed. by George L. Kumar, Lars Rudbeck.: DAKO / TRANS. from English. ed. by A. Frank and P. G. Malkov. - M., 2011. - 224 p.
- [3] Kira E. F. Evolution of reproductive surgery in gynecology and obstetrics // gynecology and obstetrics at the gynecol. - 2007. - No. 5. - P. 59-62.
- [4] Kulakov V. I. New technology and research priorities in obstetrics and obstetrics // Gynecology. - 2007. - Vol. 9, No. 1. - S. 5-7.
- [5] Strizhakov A. N., Davydov A. I., Belotserkovtseva L. D. minimally Invasive surgery in gynecology. M.: Medicine, 2001.
- [6] Kulakov V. I., Serebrennikova K. G., Volkov N. And. Chumakova N. In. Kuznetsova E. P., Menesheva V. F. Laparoscopy and hysteroscopy in programs of auxiliary reproductive technologies // Collection of materials of VI Russian forum "Mother and child". - Moscow, 2004. - S. 395-396.
- [7] Jackson T., Einarsson J. Single-port gynecologic surgery // Obstet. Gynecol. J. - 2010. - Vol. 3, N 3. - P. 133 - 39.
- [8] Mohamed ML, Nouh AA, El-Behery M. Effect on ovarian reserve of laparoscopic bipolar electrocoagulation versus laparotomic hemostatic sutures during unilateral ovarian cystectomy. Int J Gynaecol Obstet 2011; 114 (1): 69 -72.

CONTENTS

Volume 6, Number 6, December 2016

Treatment Tactics of Transcondylar and Supracondylar Fractures of Humerus Complicated by Peripheral Circulation Disturbances in Children F. A. Masharipov	167
Minimally Invasive Interventions in Surgery of Perforated Duodenal Ulcers Sh. I. Karimov, M. Sh. Khakimov, Sh. E. Ashurov	171
Ophthalmological Complications from Diabetes Mellitus: The Organization of Medical Care and Peculiarities of Disability Formation in Tashkent S. A. Djamalova, S. T. Iskandarova, A. S. Babadjanov	176
Minimally Invasive Surgery in Restoring Reproductive Function of Female Infertility Caused by Benign Ovarian Structural Changes Shukurov Farkhad Ishkulovich	182
Drug Counterfeiting: The Situation in the Gaza Strip Mohammed Abuiriban, Sami El Deeb	186
Acute Obstructive Ileus as a Late Postoperative Complication of Peptic Ulcer Disease M. Sh. Khakimov, U. K. Nasritdinov, Sh. E. Ashurov, D. Khusinov	193

ISSN: 2165-901X

