

**TOSHKENT TIBBIYOT AKADEMIYASI
«O'QUV VA ILMIY TIBBIYOT JURNALI»**

**TASHKENT MEDICAL ACADEMY
«JOURNAL OF EDUCATIONAL AND SCIENTIFIC MEDICINE»**

IXTISOSLASHUVI: «TIBBIYOT SOHASI»

(ISSN 2181-3175)

Berilgan sanasi: 31-01-2022

Mazkur hujjat Vazirlar Mahkamasining 2017 yil 15 sentabrdagi 728-son qarori bilan tasdiqlangan O'zbekiston Respublikasi Yagona interaktiv davlat xizmatlari portali to'g'risidagi nizomga muvofiq shakllantirilgan elektron hujjatning nusxasi hisoblanadi.

№ 1 (05), 2022 (Том 1)

**BOSH MUHARRIR – Shadmanov A.K.
BOSH MUHARRIR O'RINBOSARI – Oxunov A.O.**

MA'SUL KOTIB:
Boboqulova Sh.A.
TAHRIRIYAT JAMOASI:

Azizova F.L.
Alyavi A.L.
Boymurodov Sh.A.
Karimov N.F.
Karimov Sh.I.
Tuychiev L.N.
Shayxova G.I.

TAHRIRIYAT KENGASHI:
Abdullaeva R.M.
Azizova F.X.
Ahmedov X.S.
Ashurov Z.Sh.
Irisqulov B.Y.



Karimjonov I.A.
Mamatqulov B.M.
Meliboeva R.N.
Mirahmedova X.T.
Mirtazaev O.M.
Navro'zov B.S.
Norqulov S.D.
Rahimboyeva G.S.
Salomova F.I.
Tashkenbayeva U.A.
Tursunov X.Z.
Usmonxo'jayeva A.A.
Xudayqulova G.K.
Xjanazarov I.E.
Yo'ldosheva D.Yu.

Публикации в журнале учитываются как опубликованные работы при защите диссертаций на соискание ученых степеней Узбекистана и зарубежья.

К публикации принимаются статьи, ранее нигде не опубликованные и не представленные к печати в других изданиях. Статьи, поступившие в редакцию, рецензируются. За достоверность сведений, изложенных в статьях, ответственность несут авторы публикаций. При использовании материалов ссылка на журнал и авторов статей обязательна.

Материалы публикуются в авторской редакции.

Tahririyat manzili:

Tashkent Medical Academy 100109, Tashkent, Uzbekistan
Farabi street 2 Helpline: +99878-150-7825:
Fax: +998 78 1507828 Email:info@tma.uz

Khalikov S.P., Karimov M.R., Butaev L.A., Holmamatov H.T., Ahadov M.M. Acute tracheal injury	77
Abdurakhmanov F.M. Impact of COVID-19 on the course of diabetic foot syndrome.....	87
Sherali A.Kh. Immediate results of endovascular and little invasive methods of treatment of lung purulent diseases with diabetes mellitus.....	93
Matmurotov K.J. Defeat of angiosoma in purulent-necrotic processes of foot in patients with diabetic gangrene of lower limbs	97
Bobobekov A.R. Optimization of diagnostic and treatment methods acute abscesses and gangren of lungs in patients with diabetes	102
Iriskulov B.U., Tursunkhojaeva L.T. Comparative analysis of attention, memory and thought's potential of students.....	107
Okhunov A.O., Khudaibergenova N.Sh., Atakov S.S., Bobabekov A.R., Kasimov U.K. Role and place of technologies webinar in cooperation of the educational process of the branches of the tashkent medical academy	112
TursunovKh.Z., Nishanov D.A. Morphological diagnosis - goals, objectives, opportunities	118
Rajabova R.Sh., Nurillaeva N.M. Multifactor relationship of physical activity with other risk factors for coronary heart disease	125
Kasimov U. Analysis of the results of treatment of 93 patients with post-COVID thrombosis of the cavernous sines in the conditions of the department of purulent surgery.....	131
Marasulov A.F. To optimization of the independent kind of activity of students of a medical university.....	138

IMMEDIATE RESULTS OF ENDOVASCULAR AND LITTLE INVASIVE METHODS OF TREATMENT OF LUNG PURULENT DISEASES WITH DIABETES MELLITUS

Sherali Abdikhamidovich Khamdamov
Tashkent Medical Academy

РЕЗЮМЕ

Проведение длительной внутриартериальной катетерной терапии у больных с гнойно-некротическими поражениями стопы при сахарном диабете позволило значительно улучшить результаты лечения, сократить число высоких ампутаций нижних конечностей до 2,7% и избежать раннего рецидива гнойно-некротического процесса на стопе.

XULOSA

Qandli diabet fonida kechuvchi oyoqlar yiringli nekrotik jarayonlarida uzoq muddatli areriya ichi katetr terapiyasini qo'llash yiringli nekrotik jarayonlarni erta qaytalanishidan tashqari, oyoqlar yuqori amputatsiyasini 2,7% qisqartiradi.

ABSTRACT

Conducting long-term intra-arterial catheter therapy in patients with purulent-necrotic lesions of the foot in diabetes mellitus significantly improved the results of treatment, reduced the number of high amputations of the lower extremities to 2.7% and avoided early recurrence of purulent-necrotic process in the lungs.

Diabetes mellitus (DM) is one of the serious problems, the scales of which are being increased and concerned the people of all ages and of all countries. The earliest of all diseases disability, high death rate (III-rd place after cardiovascular pathologies and malignant newgrowths) have been defined diabetes mellitus as first priorities of national health systems all over the world, consolidated by Saint-Vincent declaration. The world number of patients with diabetes has been exceeded 100 mln. people. The number of patients is annually increased in 5-7%, and every 12-15 years is doubled (1,2).

One of the most frequent complications of DM is purulent-necrotic damage of lung. (3,4,5). Even mild inflammatory phenomena can have grievous consequences in these patients. There are ideal conditions for surgical infection progressing on the so-called diabetic lung, formed as a result of diabetic micro and macro angiopathy, osteoartropathy and neuropathy (1,4). Despite the advantages of up-to-date medicine, the percentage of stumps and lethality from DM combined with purulo-surgical infection are high enough and attain accordingly up to 35% till 65% and up to 6% till 44% (2,5).

The improvement of surgical treatment of purulo-necrotic lung damages (PNFD) in diabetes mellitus, diabetic gangrene has been seen in the development of methods for preservation of functioning stump. Little is known in literature about attempts of applying protracted intra-arterial infusions of various medicinal preparations in PNFD in patients with diabetes mellitus (1,4,6). But the subject remains disputable and requires the solution of the problem of the efficiency of therapeutical indication. It should be developed the techniques of its application, the possible complications aren't studied, and the ways of its prophylactics are not worked. No doubt that the compound of infusat should include medicinal preparations, affected on different pathogenetical sections of purulo-necrotic process of lung against a background of changes occurred in tissues in diabetes. The question about the regime of intra-arterial infusion of antibacterial agents is the most important there.

In a view of aforesaid, the research aim is to increase treatment efficiency and to decrease the rate of stumps in patients with PNFD in diabetes mellitus by using protracted regional intra-arterial infusion of medicinal

preparations into the complex of treatment activities.

MATERIAL AND METHODS OF INVESTIGATION

This work has been based on the analysis of examination results and treatment of

156 patients admitted to the Republican centre of contaminated surgery with diabetes mellitus (DM), complicated by purulo-necrotic lung damages (PNFD), for the period from 2006 till 2015 yy.

According to applied treatment techniques the patients were separated into three groups. The control group consisted of 29 patients. They were treated by generally accepted method and at the same time they were not given PIACT by different reasons. I basic group included 90 patients, who were administered protracted intra-arterial catheter therapy with drop-by-drop continuous introduction of antibiotics in their complex treatment. In this group, operative measures as necrectomy and amputations were made after PIACT completion and improvement of the condition of affected extremity. II basic group consisted of 37 patients, they were given protracted intra-arterial catheter therapy with periodical stream introduction of antibiotics (50% of daily dose) against a background of their drop-by-drop introduction (50% of daily dose). Operative measures as necrectomy and economical amputations were carried out after admission to a hospital or within first 2-3 days, whereupon PIACT was made to patients in order to strength the results of operation and reduce the inflammatory process in postoperative period.

The necessity of performing early necrectomy in diabetic mellitus is determined by the presence of apparent intoxication from the pathologic nidus against a background of waning in patients. Under these circumstances, the interruption of intoxication by cutting of purulo-necrotic focus allows to prevent the process of generalization. Compared groups were representative by sex, age, duration and severity of diabetic disease, concomitant diseases, scope of purulo-necrotic lung damage and duration of out-patient treatment as well.

Besides general clinical examinations of blood and urine in patients, biochemical blood composition was determined, including the

control of glycemia level. The complex of instrumental examination included Doppler ultrasonography and transcutaneous determination of oxygen tension (TcPO₂) for all patients. The patients were made angiographic studies and X-ray of lung by indications. Microbiological studies were made under the conditions of aero and anaerobiosis. The material for microbiological studies was the wound effluent and also necrotic tissues, taken directly from the wound.

STUDY RESULTS In II and III clinical groups all patients were given protracted intra-arterial catheter therapy.

When developed PIACT technique, we used catheterization of contra lateral femoral artery, directing catheter on damaged side by Beckman. Catheterization was made throughout pathologically unchanged contra lateral femoral artery. It was infused into rather big distance into the artery lumen that practically excluded the possibility of its going out from the artery lumen. The medications were part of introduced infusat and were directed on angiospasm removal and intoxication, improvement of reological blood quantities and microcirculation, antibiotics of wide use according to the sensitivity of microorganisms.

The list of main ingredients from the daily dose is below. Reopolyglukin-400ml, Haemodez-400ml, No-Spa-4,0, Trental-5,0, Heparin-5000un.

Of antibiotics, Cephalosporins were more often used: Klaphoran, Cephamsin + aminoglycosides: Gentamicin in combination with Metronidazole or Dalacin in the average therapeutical doses.

The general volume of infusat was from 800 to 1000 ml a day. The appearance of sapful granulations into the wound, reduction of intoxication, improvement of common state of patient was the evidence to stop PIACT. Microbiological studies showed that St.aureus – 36,9%, E.coli – 18,9% and Ps.aerogenosa – 18,4% occurred among aerobes most of all. They were more sensitive to Klarophan, Cephamsin, and Oxacillin.

Pr.Melaninogenicus 41,5% and B.flagilis 10,5% often occurred among anaerobes. They were more sensitive to clindamaine, Metronidazole. Having analyzed the re-

sults of treatment in all groups, they were separated into good, satisfactory and unsatisfactory. The results of treatment considered as good if PNFD in patients with diabetes completely stopped and the whole supporting function of lung was saved and work activity restored; satisfactory, if PNFD regressed after the carried-out treatment measures, the supporting function of lung was saved, but work activity was restored partly; unsatisfactory, if PNFD progressed in spite of the complex therapy and we had to ablate the extremity.

Our data proved the efficiency of infusion of protracted intra-arterial catheter therapy into the complex therapy to patients with PNFD against a background of diabetes.

Its inclusion into complex therapy allowed increasing the number of good results in 23% in I basic group and decreasing the number of bad results from 44,9% to 27,5%. i.e., if the treatment in control group finished with high ablation of lower extremity leading to total disability in 44,9% of patients, so in I basic group this rate was 27,7%.

In control group the results were estimated as good in 1 (3,4%) patient, satisfactory in 13(44,9%) patients, unsatisfactory in 13(44,9%) patients. The lethal outcome happened in 2(6,8%). Reoperations were performed on 8 patients. The average bed day was $33,7 \pm 3,45$.

In I basic group, the results were estimated as good in 32 (35,4%) patients, satisfactory in 33(37%) patients, unsatisfactory in 25(27,5%) patients, lethal outcome was in 5(5,5%) patients. Repeated operations were performed on 22 patients. The average bed-day in this group was $26,9 \pm 1,76$ ($p < 0,55$). But even these rates can be scarcely considered satisfactory; in this connection we made attempts to find the causes of unsatisfactory results in I basic group and the ways of their eliminations, as well.

Obtained analysis of the clinical dynamics, cytological and bacteriological pictures showed the little efficiency of round-the-clock drop-by-drop introduction of antibiotics. In this connection, in II basic group, we carried out the procedure of periodical against a background drop – by – drop, stream intra-arterial introduction of high doses of antibiotics. As

compared with continuous drop-by-drop introduction, the stream introduction of antibiotics due to lesser blood volume, in which the preparation is being dissolved, provides high peak concentration of active substance in nidus, depending on the time of its introduction in 7-13 times. The carried out unfolded microbiological studies of the substance separated from the wound showed the efficiency of this procedure and its use allowed decreasing the level of microbe seeding of representatives of aerobes and anaerobes flora to critical (fewer than 10^4 KOE/ml) on third day.

On this background, in II basic group we performed early economical operations on purulo-necrotic lung damage against a background of diabetic mellitus directed on the transformation of septic wound into open cleaned one under the shelter of protracted intra-arterial catheter therapy by proposed technique. Should be noticed that economical operations in this group had been performed for 24 hours before starting catheter therapy in 23 patients and on the 1-2 days of its use in 11 patients else. The use of our proposed technique in II basic group allowed improving the results of treatment significantly. In this way, the number of high amputations of low extremities in this group reduced to 2,7%, as against 27,5% and 44,9% in I basic and control groups, correspondingly. It should be specially noticed that in the II basic group we didn't register out any case of early recurrence of purulo-necrotic process on lung after PIACT.

DISCUSSION

In study process we had to be dealt with unpleasant moments connected with the occurrence of complications. The analysis of critical material showed that they generally occurred at the initial level of applied therapy. So, when analyzed complications, related with catheterization of the artery and the performance of PIACT became obvious that they developed in 12(9,4%) patients. At that, only in 4 cases (false aneurisms) of abdominal aorta-1, thrombosis of artery-1 and fallacious aneurisms of femoral artery-2 - these complications were related with catheterization of the artery. The patients were required surgical correction. 10 of 12 cases of complications were registered out in I basic group patients and 8 ones occurred in

shakedown period of technique. There were only 2 complications which were not related with catheterization in II basic group: 1 patient had a reaction on contrast and 1 had reaction on medication. This allows us considering that this procedure is more harmless and reliable.

In II basic group we estimated the results as positive in 12 (37,7%) patients, satisfactory in 24(59,8%) patients, unsatisfactory in 1(2,75) patients, fatal outcome occurred in 2 patients. Reoperations data were performed on 4 patients. The average bed-day in group was $19,2 \pm 1,69$ ($p < 0,55$).

Thus, the obtained data convincingly proved the efficiency of including PIACT into the complex of treatment for patients with PNFD against a background of diabetes, and its use in combination with early economical operations on lung by our proposed scheme, directed on conversion of purulent wound into open cleaned one, this was allowed to reduce purulo-necrotic process and preserve the supporting function of lower extremity in most of patients.

In addition to, the early performance of necrectomy ensured liquidation of nidus before developing the generalization of infectious toxic process. Subsequently, it has been noted rapid reduction of toxemia manifestation and normalization of liver and kidneys functions.

REFERENCES:

- 1) I.V. Guriyeva, I.V.Kuzina, A.V.Voronin et al., Particularities of diagnostics and treatment of diabetic lung damages // Surgery.-2005.-№ 10.-C.39-43.
- 2) A.V.Zemliyanoy, A.A.Paltsin, A.M.Svetuhin, et al., Substantiation and variants of surgical approach of complex surgery of purulo-necrotic form of "diabetic lung"//Surgery.-2003.-№ 10.-C. 44-48.
- 3) A.M. Svetuhin, A.B.Zemlyanoy, Sugecal approach to the treatment of purulo-necrotic forms of "diabetic lung" //Works of scientific and practical conferences: Modern aspects of diagnostics, treatment, prevention of lower extremity damages in patients with diabetic mellitus. – M., 2008:-C.167-177.
- 4) B.S.Briskin, E.A.Tartakovskiy, N.A.Gvozdev et al., The treatment of complications //Surgery.-2010.-№ 10.-C.53-56.
- 5) E.V.Kuleshov, Features of clinics, diagnostics and treatment of "diabetic lung" //Works of scientific and practical conferences: Modern aspects of diagnostics, treatment, and prevention of lower extremity damages in patients with diabetic mellitus. – M., 2006: - C.138-142.
- 6) S.M.Finegold, W.L.George, M.E.Milligan, Anaerobic infections, part I.//Dis Mon. -2005.-Vol 31.-PI.