

ISSN 2304–2338 (печатная версия)
ISSN 2413–4635 (электронная версия)

Проблемы современной науки и образования 2022. № 7 (176)

Российский импакт-фактор: 1,72

ИЗДАТЕЛЬСТВО
«Проблемы науки»

Журнал
зарегистрирован
Федеральной
службой по надзору
в сфере связи,
информационных
технологий и
массовых
коммуникаций
(Роскомнадзор)
Свидетельство
ПИ №ФС77– 47745

НАУЧНО-МЕТОДИЧЕСКИЙ ЖУРНАЛ

Главный редактор: Вальцев С.В.
Зам.главного редактора Кончакова И.В.

РЕДАКЦИОННЫЙ СОВЕТ:

Абдуллаев К.Н. (д-р филос. по экон., Азербайджанская Республика), *Алиева В.Р.* (канд. филос. наук, Узбекистан), *Акбулаев Н.Н.* (д-р экон. наук, Азербайджанская Республика), *Аликулов С.Р.* (д-р техн. наук, Узбекистан), *Ананьева Е.П.* (д-р филос. наук, Украина), *Асатурова А.В.* (канд. мед. наук, Россия), *Аскарходжаев Н.А.* (канд. биол. наук, Узбекистан), *Байтасов Р.Р.* (канд. с.-х. наук, Белоруссия), *Бакико И.В.* (канд. наук по физ. воспитанию и спорту, Украина), *Бахор Т.А.* (канд. филол. наук, Россия), *Баулина М.В.* (канд. пед. наук, Россия), *Блейх Н.О.* (д-р ист. наук, канд. пед. наук, Россия), *Боброва Н.А.* (д-р юрид. наук, Россия), *Богомолов А.В.* (канд. техн. наук, Россия), *Бородай В.А.* (д-р социол. наук, Россия), *Волков А.Ю.* (д-р экон. наук, Россия), *Гавриленкова И.В.* (канд. пед. наук, Россия), *Гарагонич В.В.* (д-р ист. наук, Украина), *Глуценко А.Г.* (д-р физ.-мат. наук, Россия), *Грищенко В.А.* (канд. техн. наук, Россия), *Губарева Т.И.* (канд. юрид. наук, Россия), *Гутникова А.В.* (канд. филол. наук, Украина), *Датий А.В.* (д-р мед. наук, Россия), *Демчук Н.И.* (канд. экон. наук, Украина), *Дивненко О.В.* (канд. пед. наук, Россия), *Дмитриева О.А.* (д-р филол. наук, Россия), *Доленко Г.Н.* (д-р хим. наук, Россия), *Есенова К.У.* (д-р филол. наук, Казахстан), *Жамулдинов В.Н.* (канд. юрид. наук, Казахстан), *Жолдошев С.Т.* (д-р мед. наук, Кыргызская Республика), *Зеленков М.Ю.* (д-р полит. наук, канд. воен. наук, Россия), *Ибадов Р.М.* (д-р физ.-мат. наук, Узбекистан), *Ильинских Н.Н.* (д-р биол. наук, Россия), *Кайракбаев А.К.* (канд. физ.-мат. наук, Казахстан), *Кафтаева М.В.* (д-р техн. наук, Россия), *Кикаидзе И.Д.* (д-р филол. наук, Грузия), *Клишков Г.Т.* (PhD in Pedagogic Sc., Болгария), *Коблано Ж.Т.* (канд. филол. наук, Казахстан), *Ковалёв М.Н.* (канд. экон. наук, Белоруссия), *Кравцова Т.М.* (канд. психол. наук, Казахстан), *Кузьмин С.Б.* (д-р геогр. наук, Россия), *Куликова Э.Г.* (д-р филол. наук, Россия), *Курманбаева М.С.* (д-р биол. наук, Казахстан), *Курпаяниди К.И.* (канд. экон. наук, Узбекистан), *Линькова-Даниельс Н.А.* (канд. пед. наук, Австралия), *Лукиенко Л.В.* (д-р техн. наук, Россия), *Макаров А. Н.* (д-р филол. наук, Россия), *Мацаренко Т.Н.* (канд. пед. наук, Россия), *Мейманов Б.К.* (д-р экон. наук, Кыргызская Республика), *Мурадов Ш.О.* (д-р техн. наук, Узбекистан), *Мусаев Ф.А.* (д-р филос. наук, Узбекистан), *Набиев А.А.* (д-р наук по геоинформ., Азербайджанская Республика), *Назаров Р.Р.* (канд. филос. наук, Узбекистан), *Наумов В. А.* (д-р техн. наук, Россия), *Овчинников Ю.Д.* (канд. техн. наук, Россия), *Петров В.О.* (д-р искусствоведения, Россия), *Радкевич М.В.* (д-р техн. наук, Узбекистан), *Рахимбеков С.М.* (д-р техн. наук, Казахстан), *Розьходжаева Г.А.* (д-р мед. наук, Узбекистан), *Романенкова Ю.В.* (д-р искусствоведения, Украина), *Рубцова М.В.* (д-р социол. наук, Россия), *Румянцев Д.Е.* (д-р биол. наук, Россия), *Самков А. В.* (д-р техн. наук, Россия), *Саньков П.Н.* (канд. техн. наук, Украина), *Селитренникова Т.А.* (д-р пед. наук, Россия), *Сибирцев В.А.* (д-р экон. наук, Россия), *Скрипка Т.А.* (д-р экон. наук, Украина), *Солов А.В.* (д-р ист. наук, Россия), *Стрекалов В.Н.* (д-р физ.-мат. наук, Россия), *Стукаленко Н.М.* (д-р пед. наук, Казахстан), *Субачев Ю.В.* (канд. техн. наук, Россия), *Сулейманов С.Ф.* (канд. мед. наук, Узбекистан), *Трегуб И.В.* (д-р экон. наук, канд. техн. наук, Россия), *Уноров И.В.* (канд. юрид. наук, д-р ист. наук, Россия), *Федоскина Л.А.* (канд. экон. наук, Россия), *Хилтухина Е.Г.* (д-р филос. наук, Россия), *Цурцян С.В.* (канд. экон. наук, Республика Армения), *Чиладзе Г.Б.* (д-р юрид. наук, Грузия), *Шамишина И.Г.* (канд. пед. наук, Россия), *Шарипов М.С.* (канд. техн. наук, Узбекистан), *Шевко Д.Г.* (канд. техн. наук, Россия).

Издается с 2011
года

Территория
распространения:
зарубежные
страны,
Российская
Федерация

Подписано в
печать:

04.10.2022.

Дата выхода в
свет:

06.10.2022

Формат 70x100/16.

Бумага офсетная.

Гарнитура

«Таймс».

Печать офсетная.

Усл. печ. л.

Тираж 1 000 экз.

Заказ №

Свободная цена

© ЖУРНАЛ «ПРОБЛЕМЫ СОВРЕМЕННОЙ НАУКИ
И ОБРАЗОВАНИЯ»/PROBLEMS OF MODERN SCIENCE
AND EDUCATION»

© ИЗДАТЕЛЬСТВО «ПРОБЛЕМЫ НАУКИ»

МЕДИЦИНСКИЕ НАУКИ..... 29

Abdumutalipov U.Sh., Kosimov K.K. LONG-TERM RESULTS OF ENDOSCOPIC OPERATIONS IN CHRONIC SINUSITIS / Абдумуталипов У.Ш., Косимов К.К. ОТДАЛЕННЫЕ РЕЗУЛЬТАТЫ ЭНДОСКОПИЧЕСКИХ ОПЕРАЦИЙ ПРИ ХРОНИЧЕСКИХ СИНУСИТАХ..... 29

Saliahunova Kh.O., Shaykhova H.E. TO INCREASE THE EFFECTIVENESS OF MODERN METHODS OF TREATMENT OF PATIENTS WITH CHRONIC OBSTRUCTIVE PULMONARY DISEASE AND THE OBSERVED CONCOMITANT MANIFESTATIONS OF RHINOSINUSITIS / Саляхунова Х.О., Шайхова Х.Э. ПОВЫСИТЬ ЭФФЕКТИВНОСТЬ СОВРЕМЕННЫХ МЕТОДОВ ЛЕЧЕНИЯ ПАЦИЕНТОВ С ХРОНИЧЕСКОЙ ОБСТРУКТИВНОЙ БОЛЕЗНЬЮ ЛЕГКИХ И НАБЛЮДАЕМЫМИ СОПУТСТВУЮЩИМИ ПРОЯВЛЕНИЯМИ РИНОСИНУСИТА..... 32

Ulmasov A.O., Kosimov K.K. IMMUNOHISTOCHEMICAL ASSESSMENT OF SIGNS OF INFLAMMATION OF THE NASAL CAVITY AND PARANASAL SINUSES IN ALLERGIC RHINITIS / Улмасов А.О., Косимов К.К. ИММУНОГИСТОХИМИЧЕСКАЯ ОЦЕНКА ПРИЗНАКОВ ВОСПАЛЕНИЙ ПОЛОСТИ НОСА И ОКОЛОНОСОВЫХ ПАЗУХ ПРИ АЛЛЕРГИЧЕСКОМ РИНИТЕ..... 36

Ulmasov B.B., Arifov S.S. THE MECHANISM OF DEVELOPMENT AND IMPROVEMENT OF TREATMENT OF ACUTE SINUSITIS OCCURRING WITH A BLOCK OF NATURAL ANASTOMOSIS / Улмасов Б.Б., Арифов С.С. МЕХАНИЗМ РАЗВИТИЯ И СОВЕРШЕНСТВОВАНИЕ ЛЕЧЕНИЯ ОСТРЫХ СИНУСИТОВ, ПРОТЕКАЮЩИХ БЛОКОМ ЕСТЕСТВЕННОГО СОУСТЬЯ..... 39

ИСКУССТВОВЕДЕНИЕ 44

Rakhimov B.M. СТИЛИ СОПРОВОЖДЕНИЯ ГОЛОСОМ И СОТВОРЧЕСТВА В ХОРЕЗМСКОМ ПЕСЕННОМ ИСКУССТВЕ / Rakhimov B.M. STYLES OF VOICE ACCOMPANIMENT AND CO-CREATION IN THE KHOREZM SONG ART 44

Nesterova A.S., Davydova T.Yu. ЗАРОЖДЕНИЕ И СТАНОВЛЕНИЕ ОПЕРНОГО ИСКУССТВА В ЕВРОПЕЙСКОЙ И УЗБЕКСКОЙ КУЛЬТУРАХ / Nesterova A.S., Davydova T.Yu. THE ORIGIN AND FORMATION OF OPERA ART IN EUROPEAN AND UZBEK CULTURES 47

Allanbayev R.O. ИСПОЛНИТЕЛЬСКОЕ ИСКУССТВО КАРАКАЛПАКСКОГО ЭПОСА: ИСТОРИЯ, РАЗВИТИЕ / Allanbayev R.O. PERFORMING ART OF THE KARAKALPAK EPIC: HISTORY, DEVELOPMENT 50

Abdugapparov A.A. НАСЛЕДИЕ ВЕЛИКИХ МЫСЛИТЕЛЕЙ ВОСТОКА – ФУНДАМЕНТ МУЗЫКАЛЬНО-ПЕДАГОГИЧЕСКОЙ КОНЦЕПЦИИ ПРОФЕССОРА РАУФА КАДЫРОВА / Abdugapparov A.A. THE LEGACY OF THE GREAT THINKERS OF THE EAST IS THE FOUNDATION OF THE MUSICAL AND PEDAGOGICAL CONCEPT OF PROFESSOR RAUF KADYROV..... 53

Abdugapparov A.A. ИСТОЧНИКИ ТВОРЧЕСТВА / Abdugapparov A.A. SOURCES OF CREATIVITY..... 56

ПОВЫСИТЬ ЭФФЕКТИВНОСТЬ СОВРЕМЕННЫХ МЕТОДОВ ЛЕЧЕНИЯ ПАЦИЕНТОВ С ХРОНИЧЕСКОЙ ОБСТРУКТИВНОЙ БОЛЕЗНЬЮ ЛЕГКИХ И НАБЛЮДАЕМЫМИ СОПУТСТВУЮЩИМИ ПРОЯВЛЕНИЯМИ РИНОСИНУСИТА

Саляхунова Х.О.¹, Шайхова Х.Э.²

¹Саляхунова Хуришидахон Одиловна – ассистент;

²Шайхова Холида Эркиновна – доктор медицинских наук, профессор,
кафедра оториноларингологии,
Ташкентская медицинская академия,
г. Ташкент, Республика Узбекистан

Аннотация: коморбидный фон пациентов с хронической обструктивной болезнью легких (ХОБЛ) отягощен отнюдь не меньше соматического статуса «сосудистых» больных, при этом очевидно, что ХОБЛ в свою очередь по ряду клинических и лабораторных показателей усугубляет клиническое течение абсолютного большинства известных сегодня заболеваний.

В этой статье обсуждается идея хронической обструктивной болезни легких и сопутствующих риносинуситов, современные методы их диагностики и лечения.

Болезни органов дыхания до настоящего времени представляют собой важную социально-медицинскую проблему во всем мире, поскольку по удельному весу в общей смертности населения занимают одно из ведущих мест, а экономический ущерб, наносимый обществу вследствие высокой заболеваемости и инвалидизации больных, огромен. В течение последних 25 лет общая заболеваемость болезнями органов дыхания неуклонно возрастает. На долю органов дыхания, по данным официальной статистики, приходится около 40% всех случаев заболеваемости, которая превосходит уровни заболеваемости другими классами болезней.

Ключевые слова: риносинусит, коморбидность, хронической обструктивной болезнью легких.

UDC 616.03

Relevance. Respiratory diseases are still an important socio-medical problem throughout the world, since they occupy one of the leading places in terms of their share in the total mortality of the population, and the economic damage caused to society due to high morbidity and disability of patients is enormous. During the last 25 years, the overall incidence of respiratory diseases has been steadily increasing [2, 6]. According to official statistics, the respiratory organs account for about 40% of all cases of morbidity, which exceeds the levels of morbidity in other classes of diseases [5]. In the structure of the reasons for applying for medical care, their share in various territories ranges from 29.2 to 43.5% among adults and from 65.4 to 83.8% among children.

The most common forms of respiratory diseases are acute diseases: acute respiratory viral infections, acute bronchitis and pneumonia, respectively 94.2; 4.0; 1.8%, among the adult population and 96.6; 2.9; 0.5% - among the child population [1, 4].

Currently, more than 1 billion people of all ages in all countries of the world suffer from chronic respiratory diseases (CRD). The burden of preventable CRD has a great negative impact on the quality of life and work capacity of people affected by CRD diseases that cause early death. According to WHO data obtained from a survey of representative samples of the population based on an international questionnaire and spirometry as part of the GARD (Global Alliance against Chronic Respiratory Diseases) epidemiological study, chronic respiratory diseases are a serious health problem in all countries of the world [3, 7].

Under the general name "chronic respiratory diseases" a number of serious diseases are grouped together. Preventable CRDs include asthma and respiratory allergies, chronic

obstructive pulmonary disease, occupational lung disease, and pulmonary hypertension. They pose a serious threat to society, especially among socially vulnerable groups. The forecast of WHO experts until 2020 indicates that chronic lung diseases will not only become one of the most common forms of human pathology, but will also be among the leading causes of death [4, 8].

Currently, signs of chronic damage to the broncho-pulmonary apparatus are detected in 30% of those who applied to the clinic.

Respiratory diseases are socially conditioned. The emergence of many of them is associated with the influence of various socio-hygienic factors, among which the most important are professional, environmental, social [3, 5].

Currently, from 16 to 30 main risk factors for chronic diseases of the respiratory system are distinguished, combined into several groups (social, socio-psychological, biological, environmental, socio-demographic, etc.), among which the most recognized are: dust and gas contamination of atmospheric air, professional agents, smoking (duration and intensity), alcohol abuse, repeated acute respiratory infections and acute pneumonia, unfavorable natural and climatic conditions, physical and neuropsychic overstrain, sensitization to allergens, ethnic characteristics, low socioeconomic status, aggravated premorbid background, diseases mothers during pregnancy, physical inactivity, unhealthy diet, decreased immunological reactivity [6].

The medical statistics of recent years convincingly indicates an increase in the number of diseases of the nose and paranasal sinuses, and both the overall incidence of rhinosinusitis and the proportion of this pathology in the structure of ENT diseases are increasing.

The close anatomical and physiological interrelationships of the upper and lower respiratory tract are the reason that the increase in the number of rhinosinusitis keeps pace with the increase in the number of pulmonary diseases, in particular chronic obstructive pulmonary disease (COPD), and this trend has not yet been broken, despite on the joint efforts of leading experts from around the world. COPD is the 4th leading cause of death in the over 45 age group in the world, and the number of deaths from the disease continues to increase.

It has been proven that chronic inflammation of the upper and lower respiratory tract has a similar etiology, due to the fact that the mucous membrane of the airways from the nasal cavity to the middle bronchi has a similar anatomical and histological structure [9].

It is known that one of the factors contributing to the exacerbation of COPD, and in some cases its development, is a bacterial infection [3]. This circumstance dictates the need to obtain convincing evidence of the relationship between the nature of chronic bacterial infection of the upper respiratory tract (bacterial chronic rhinosinusitis) and the severity, nature, clinical features of COPD, as well as the number of its exacerbations.

Purpose of the study. The purpose of this study is to determine the therapeutic efficacy of modern drugs in the treatment of the Complex in patients with chronic rhinosinusitis and chronic obstructive pulmonary disease.

Materials and research methods. To solve the tasks set, 40 patients were examined. All patients were divided into 2 groups of 20 people. The first group consisted of 12 men and 8 women aged 39 to 66 years (mean age 52 years). All patients of this group suffered from exacerbation of chronic rhinosinusitis in combination with COPD of the 1st-2nd stage during the exacerbation.

Research results. All patients, along with standard otorhinolaryngological and radiological examinations, underwent a bacteriological examination of nasal discharge, ultrasound (US) scanning of the maxillary sinuses, a study of mucociliary function using a saccharin test, and analyzed the results of rhinopneumomanometry. To clarify the dynamic state of the bronchopulmonary system, patients underwent a pulmonological examination (consultation of a pulmonologist, determination of respiratory function). All patients of this group were added to the treatment regimen Erespal at a dose of 160 mg per day with 2 doses. Patients in this group were also prescribed bronchodilators.

Group 2 included 10 men and 10 women aged 40 to 68 years (mean age 54 years) with exacerbation of chronic rhinosinusitis in combination with stage 1-2 COPD during the exacerbation period. The examination was carried out according to the scheme indicated above. Treatment of patients in this group consisted of the appointment of mucolytics and bronchodilators. Hormone therapy was not performed, except for patients with concomitant bronchial asthma. These patients were further excluded from the study.

We took into account such indicators as headache, malaise, difficulty in nasal breathing, nasal discharge, shortness of breath (the severity of shortness of breath was assessed according to the original Borg scale), as well as cough, the number of dry wheezing, and sputum production. The effectiveness of the drug was evidenced by the dynamics of the parameters of ultrasound scanning of the paranasal sinuses.

The results of this study were interpreted as follows: 2-4 LEDs - normal, 4-6 - I degree of edema of the mucous membrane of the maxillary sinus, 6-8 - II degree, 8-10 - III degree, gap between the LEDs - the presence of pathological content, which produced using a SIHUSCAN one-dimensional ultrasound machine with a built-in LED indicator, as well as the dynamics of mucociliary clearance and rhinopneumometry parameters. The results of the study were recorded in a specially designed card filled in by patients during the study. The severity of symptoms was assessed in points from 0 to 5 using a special scale.

Our study indicates a fairly high therapeutic efficacy of fenspiride.

In the 1st group, by the end of treatment (day 10), there was no discharge from the nose in 20 (100%).

In patients of the 2nd group, by the end of treatment (10th day), there was no discharge in 7 (35%), in 7 - the amount of discharge decreased sharply. In 6 (30%) patients, there were no positive dynamics during the treatment. In general, a positive clinical effect was observed in 14 (70%) of 20 patients.

There were no changes in the microbial landscape under the influence of fenspiride.

After analyzing the dynamics of indicators of mucociliary transport, we can conclude that while taking Erespal, the function of the ciliated epithelium improves.

In the study of respiratory function, we determined volume and speed indicators: vital capacity (VC), forced vital capacity (FVC), forced expiratory volume in 1 second (FEV1, modified Tiffno index ($FEV1 / FVC$). The obtained indicators were evaluated as a percentage relation to the norm.

The results of the study of respiratory function during the exacerbation of COPD showed that against the background of the use of fenspiride, there was an increase in FEV1 by 1.32 times (in the comparison group - 1.03 times) and there was a tendency to increase the modified Tiffno index, however, this indicator did not reach the degree of statistical significance.

From the above data, it can be concluded that the clinical signs of exacerbation of chronic rhinosinusitis and concomitant COPD both in the first and second groups of patients were significantly reduced, however, while taking fenspiride, the positive effect arose faster and more actively. In none of the cases, we noted undesirable side effects during treatment with the drug. This indicates the safety of the drug in clinical practice.

The results obtained indicate a sufficiently high therapeutic efficacy of fenspiride in relation to acute and exacerbation of chronic sinusitis with comorbid COPD, and we consider it expedient to widely use this drug in the complex treatment of these patients.

Conclusion. The results obtained indicate a sufficiently high therapeutic efficacy of fenspiride in relation to acute and exacerbation of chronic sinusitis with comorbid COPD, and we consider it expedient to widely use this drug in the complex treatment of these patients.

Thus, the assessment of comorbidity in patients with COPD is an important component of clinical prognosis. Targeted identification of comorbidity would allow taking into account the prevalence of diseases of the cardiovascular system in patients with COPD when choosing cardiosafe drugs to prolong the quality of life of patients.

References / Список литературы

1. *Bugaenko V.V., Slobodskoy B.C., Tovstukha V.V.* Comorbid conditions: IHD and chronic obstructive pulmonary disease, 2011. P. 25-27.
2. *Vertkin A.L., Skotnikov A.S., Gubzhokova O.M.* "Comorbidity in chronic obstructive pulmonary disease: the role of chronic systemic inflammation and the clinical and morphological niches of roflumilast". Attending physician 8/6. Moscow, 2014. P. 1-5.
3. *Davydovskaya E.I.* Comorbidity in chronic obstructive pulmonary disease. breast cancer. Moscow, 2013. P. 30-40.
4. *Mukatova A.M., Idrisov U.A. etc.* Cardiovascular pathology in COPD. Therapeutic Readings. Proceedings of the republican scientific-practical conference. Almaty, 2008. P. 57-60.
5. *Chereshnev V.A., Gusev E.Yu., Yurchenko H.H.* Systemic inflammation - myth or reality // Bulletin of the Russian Academy of Sciences, 2004. Vol. 74. № 3. Pp. 219-227.
6. *Chuchalin A.G., Ovcharenko S.I., Avdeev S.N.* Place of Onbrez Breeze-haler (indacaterol) in the treatment of chronic obstructive pulmonary disease: conclusion of the expert council // Pulmonology, 2011. № 6. Pp. 124-125.
7. *Pawankar R.* Allergic rhinitis and its impact on asthma: an evidence-based treatment strategy for allergic rhinitis // Asian. Pac. J. Allergy Immunol., 2002. № 3. P. 43-52.
8. *Ural A., Oktemer T.K., Kizil Y.* Impact of isotonic and hypertonic saline solutions on mucociliary activity in various nasal pathologies: clinical study. // J Laryngol Otol., 2009. Vol. 123(5). P. 517-521.
9. *Zawisza E.* Effectiveness and tolerance of fenspiride treatment in chronic sinusitis. Results of the Polish multicenter study. // Otolaryngol Pol., 2005. Vol. 59(1). P. 141-145.

IMMUNOHISTOCHEMICAL ASSESSMENT OF SIGNS OF INFLAMMATION OF THE NASAL CAVITY AND PARANASAL SINUSES IN ALLERGIC RHINITIS

Ulmasov A.O.¹, Kosimov K.K.²

Email: Ulmasov17176@scientifictext.ru

¹*Ulmasov Alijon Obidovich – Assistant;*

²*Kosimov Kobil Kosimovich - Doctor of Medical Sciences, Professor, Head of the Department, DEPARTMENT OF OTORHINOLARYNGOLOGY, ANDIJAN STATE MEDICAL INSTITUTE, ANDIJAN, REPUBLIC OF UZBEKISTAN*

Abstract: *in recent decades, allergy has become one of the most urgent medical and social problems due to the global prevalence and intensive growth of allergic diseases. Allergy occupies a leading place among other forms of pathology according to criteria such as prevalence, severity of course, complexity of diagnosis and therapy, rehabilitation, treatment costs, etc.*

Keywords: *nasal cavities, immunology, chemistry, paranasal sinuses, allergic rhinitis.*

НАУЧНОЕ ИЗДАНИЕ

ИЗДАТЕЛЬСТВО
«ПРОБЛЕМЫ НАУКИ»

АДРЕС РЕДАКЦИИ:
153008, РФ, Г. ИВАНОВО, УЛ. ЛЕЖНЕВСКАЯ, Д. 55, 4 ЭТАЖ
ТЕЛ.: +7(915)814-09-51

HTTP://WWW.IPI1.RU
E-MAIL: INFO@P8N.RU

ТИПОГРАФИЯ:
ООО «ПРЕССТО».
153025, Г. ИВАНОВО, УЛ. ДЗЕРЖИНСКОГО, Д. 39, СТРОЕНИЕ 8

ИЗДАТЕЛЬ:
ООО «ОЛИМП»
УЧРЕДИТЕЛИ: ВАЛЬЦЕВ СЕРГЕЙ ВИТАЛЬЕВИЧ;
ВОРОБЬЕВ АЛЕКСАНДР ВИКТОРОВИЧ
108814, Г. МОСКВА, УЛ. ПЕТРА ВЯЗЕМСКОГО 11/2