

This journal had been publishing since 2018

4
2022



MINISTRY OF HIGHER AND SECONDARY SPECIALIZED
EDUCATION OF THE REPUBLIC OF UZBEKISTAN

2030

UZBEKISTAN RESEARCH ONLINE

**Ministry of Health of the Republic of Uzbekistan
Tashkent Medical Academy**



Central Asian Journal of Medicine

№ 4 / 2022

Tashkent



1083

certificate

EDITORIAL BOARD

Editor-in-Chief

Alisher K. Shadmanov, DSc, M.D., Tashkent Medical Academy

Deputy Chief Editor

Oktyabr R. Teshayev, DSc, M.D., Tashkent Medical Academy

Executive Editor

Feruza Khidoyatova, DSc, M.D., Tashkent Medical Academy

Editorial Team

Anis L. Alyavi, Academician, DSc, M.D., The Republican Specialized Scientific and Practical Medical Center for Therapy and Medical Rehabilitation, Uzbekistan

Feruz G. Nazirov, Academician, DSc, M.D., The Republican Specialized Center of Surgery named after Academician V.Vakhidov, Uzbekistan

Ravshan D. Kurbanov, Academician, DSc, M.D., The Republican Specialized Cardiology Center, Uzbekistan

Shavkat I. Karimov, Academician, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Laziz N. Tuychiev, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Abdugaffar G. Gadaev, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Dilbar K. Najmutdinova, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Jae Wook Choi, DSc, M.D., Korea University, South Korea

Farkhad O. Akilov, DSc, M.D., Republican Specialized Center of Urology, Uzbekistan

Bakhtiyor U. Iriskulov, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Nargiza N. Parpieva, DSc, M.D., Republican Specialized Scientific and Practical Medical Center Of Tuberculosis And Pulmonology, Uzbekistan

Rikhsi A. Sabirova, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Feruza L. Azizova, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Barno T. Kholmatova, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Raisa I. Shalina, DSc, M.D., Department of Obstetrics and Gynecology, Pediatric Faculty, Russian National Research Medical University named after N.I. Pirogov, Russian Federation

Antanas Vaitkus, DSc, M.D., Hospital of Lithuanian University of Health Sciences Kaunas Clinics Academy of Medicine, Lithuania

Briko I. Nikolay, DSc, M.D., Honored Scientist of the Russian Federation, Academician of the Russian Academy of Sciences, Professor, Chief Epidemiologist of the Ministry of Health of the Russian Federation, Head of the Department of Epidemiology and Evidence-Based Medicine, Sechenov University, Russian Federation

Deniz Gokalp, Head of obstetrics-gynecology, reproductology, Istinye University, Turkey

Natalya Mitkovskaya, Head of Cardiology and Internal Medicine, Belarusian State Medical University, Belarusia

Utkus Algirdas, Head of Human and Medical Genetics, Vilnius University, Lithuania

Jin-San Zhang, Professor, Pharmaceutical Sciences, Wenzhou Medical University, China

Andreas Unterberg, Professor, Neurosurgery, University of Heidelberg, Germany

Maya Dgebuadze, Professor, Normal human anatomy, Tbilian republican medical university, Georgia

Yulia Stepanova, Professor, Surgery and Surgical Technologies, Moscow State University of Medicine and Dentistry named after A.I. Evdokimov, Russia

Jose Maria Peinado Herreros, Professor, Biochemistry and Molecular Biology III and Immunology, Granada University, Spain

Haris Khwaja, Professor, Clinical Lead & Consultant Bariatric Surgeon, Chelsea and Westminster Hospital, England

Daniel Truong, Professor, Neurology, UC Riverside, USA

Tomas Xolzenbayn, Professor, Surgery, Medical University of Salzburg, Austria

Ahsan Nazeer, Assistant professor, Child and adolescent psychiatry, Michigan State University College of Human Medicine, USA

Mustafa Taskin, Professor, Surgery, Istanbul University, Turkey

Galina V. Yarovenko, DSc, M.D., Professor of Surgery Department, Samara State Medical University, Russian Federation

Alisher O. Okhunov, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Nargiza M. Nurillaeva, DSc, M.D., Tashkent Medical Academy, Uzbekistan

Assistant Editor

Abror Musakhonov, Tashkent Medical Academy, Uzbekistan

Jakhongir Khamdamov, Tashkent Medical Academy, Uzbekistan

Journal edited and printed in the computer of
Tashkent Medical Academy editorial department

Editorial board of Tashkent Medical Academy

Head of the department: M.N. Aslonov

English language editor: A.Kh. Juraev

Designer corrector: S.R. Sultonova

Organizer: Tashkent Medical Academy

eISSN 2181-1326



9 772181 132009

INTRAOPERATIVE REASONS FOR CONVERSION OF LAPAROSCOPIC CHOLECYSTECTOMY TO OPEN SURGERY (SYSTEMATIC REVIEW)

**Khakimov M.Sh.¹, Karimov R.A.¹, Jasmin Sabanovic², Karim Belhaj²,
Bijendra Patel²**

¹Department of Faculty and Hospital Surgery No. 1
of the Tashkent Medical Academy, Uzbekistan

²Barts Cancer Institute, Queen Mary, University of London,
Charterhouse Square, London, UK

ABSTRACT

Three electronic databases were searched to identify the studies reporting intraoperative risk factors of conversion in laparoscopic cholecystectomy. Primary outcomes were to spot and specify the unpredictable risk factors and identify the most common intraoperative conversion causes. Secondary outcomes were comparing the impact of each group reasons on conversion rate – patient-related factors, surgeon-related factors and equipment-related and other factors of conversion.

Key words: Laparoscopic cholecystectomy, conversion, intraoperative risk factors.

INTRODUCTION

The reasons for conversion have been investigated deeply to reduce the rate of conversion. Authors of many studies in this field mainly focused on the risk factors that can be identified and diagnosed before surgery to prevent the risk of conversion. Advanced age, male gender, high body mass index (BMI), previous upper abdominal surgery, choledocholithiasis, urgent surgery, thicker gallbladder wall, raised white cell count (WCC), and alkaline phosphatase (ALP) was found as a predictor of conversion in laparoscopic cholecystectomy (A.S.Y. Hu et al, 2017 and J.P. Rothman et al, 2016).

There is a clear lack of investigations in the literature for unpredictable causes of conversion in minimally access surgery. Authors of some studies have mentioned some unpredictable risk factors (A. Gangemi et al, 2017 and A.

Licciardello et al, 2014) but they did ignore their importance in practice. Deep knowledge of the epidemiology of intraoperative risk factors for conversion to open surgery allows early identification and recognition of difficult situations and potential reasons for conversion. This gives an opportunity of avoiding unnecessary laparoscopic attempts that increases the risk of further complications that cannot be solved even open approach afterward. Moreover, keeping in mind these factors improves the situation awareness of surgeons and forces them to rethink before making any important step of the procedure that leads to minimization of severe iatrogenic complications. Further, a systematic assessment of such factors improves patients' consent forms for laparoscopic cholecystectomy and allows them to be better informed of the risks before the surgery.

Purpose of the study. The study's main aim was to define the intraoperative reasons for conversion from laparoscopic cholecystectomy to open surgery. Moreover, we aimed to reveal the popularity of each reason among others calculating their proportion within the included studies.

METHODS

Protocol and registration. This review was conducted according to the guidelines set by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) (D. Moher et al, 2009). Before analyzing the factors under consideration within the included studies and data extraction stage, the review was registered on the International Prospective Register of Systematic Reviews (PROSPERO) database (University of York, 2019) with the registration number CRD42019135834.

Eligibility criteria. We found the articles eligible for our study that reported intraoperative reasons or all causes of conversion (where intraoperative factors were mentioned as well) from standard 4-port laparoscopic cholecystectomy to open surgery. In all included studies, indication for surgery had to be gallstone disease and other benign pathologies of gallbladder. Only English language studies were included in the review.


Search strategy and study selection. A systematic search of relevant articles was carried out through the electronic medical literature databases of PubMed, Cochrane, and EMBASE by the first two authors. The search was started on January 21 2019, and refreshed on weekly basis until July 1, 2019. Titles and abstracts of the papers retrieved applying the search tactics mentioned above and those from additional databases were reviewed independently by two study authors (R.K. and J.S.) to identify studies that potentially meet the eligibility criteria. Duplicate papers in all three databases were matched and removed. As soon as a title or abstract of any record fit our eligibility criteria, full texts of the potentially

eligible studies were acquired and evaluated by two review team members in terms of eligibility for the systematic review. Any discrepancies and disagreements were resolved through discussion.

Data collection and data items. A unique modified data extraction form was designed by the first author and employed to systematically collect data from all eligible studies. The form was adjusted for our primary as well as secondary outcomes.

Statistical Analysis. Data were assembled in Microsoft Excel (Microsoft Corporation, Redmond, WA, USA). The overall proportion of conversion cases in the eligible studies was calculated using simple mathematic methods to learn the general conversion rate individually. All statistical analyses were accomplished on IBM SPSS Statistics Version 26 (IBM Corporation, Endicott, NY, USA). The total percentage, median, mean with standard deviation (SD), minimum, maximum, and p-value were calculated for each risk factor with an α level of 0.05. In this way, we found out the popularity of each intra-operational risk factor.

RESULTS. Overall, 1050 records were found in all medical electronic databases (please see the PRISMA 2009 Flow Diagram below). One study was identified in the reference list as eligible for the current review. We compared the records within three electronic databases by screening the titles and authors and found 302 identical studies, which were eliminated from the study. Next, 179 records were excluded, 126 of them were out of the topic, two studies involved paediatric content. 24 records were secondary studies and 27 findings were reporting isolated preoperative risk factors to create a prognostic program for safe cholecystectomy. Both types of records were removed from the review. Finally, sixty-eight were derived and read in full text by the first two authors of the study, and 46 papers were subsequently excluded owing to the following reasons: 28 studies ignored intra-operative reasons for conversion from laparoscopic cholecystectomy to open surgery, 13 records were unavailable in full text, four papers due to the language and the last one record was a comment letter for another study. Due to the difference in the style, methods, and outcomes of the studies (heterogeneity) found throughout the databases, we could not perform meta-analysis.



Graph 1. PRISMA 2009 Flow Diagram

Study characteristics

This systematic review included 22 prospective and retrospective observational studies thcovereding 40 315 patients in total (table 1).

Table 1.

Patients’ demographic parameters in the studies


Included studies	Sample size	Main age	Gender distribution	
			Male	Female
El Nakeeb et al	3269	N/A	933 (28.5%)	2336 (71.5%)
Gangemi et al	960	N/A	244 (25.4%)	716 (74.6%)
Licciardello et al	414	51.7±16.4	169 (40.8%)	245 (59.2%)
Taki-Eldin & Badawy	492	49.35±8.68	106 (21.5%)	386 (78.5%)
Malik	936	39.88±8.66	97 (10.36%)	839 (89.64%)
Simopoulos et al	1804	52.66±14.6	425 (23.6%)	1379 (76.4%)
		6		
Lo et al	70	59.9	28 (40%)	42 (60%)
Shamiyeh et al	5049	N/A	N/A	N/A
Yajima et al	407	N/A	183 (45%)	224 (55%)
Ishizaki et al	1339	55±13	600 (50.9%)	579 (49.1%)

Johanning et al	3247	N/A	N/A	N/A
Lim et al	201	N/A	82 (40.8%)	119 (59.2%)
Ercan et al	2015	Lap – 52.4±13.9* Con – 57.8±13.0*	644 (31.96%)	1371 (68.04%)
Kama et al	1000	43.8	196 (19.6%)	804 (80.4%)
Pavlidis et al	1263	54±15.15	354 (28%)	909 (72%)
Kala et al	8347	38.2	2825 (33.85%)	5522 (66.15%)
Shamim et al	1238	41.25±12	158 (12.76%)	1080 (87.24%)
Hadad et al	1385	N/A	N/A	N/A
Rashid et al	300	42.69	38 (12.67%)	262 (87.33%)
Genc et al	5164	N/A	1570 (30.4%)	3594 (69.6%)
Zhang et al	1265	N/A	430 (34%)	835 (66%)
Sikora et al	150	39±12	36 (24%)	114 (76%)


TOTAL 40 315

*Ercan et al calculated the mean age of the patients separately in two different groups, successful laparoscopic and converted groups

Most of the studies remain under 10 percent in terms of conversion rate (graph 2). This pattern is exceeding a quarter in Lim et al, 2007 and reached up to 20 percent in the study conducted by Sikora et al. (1995). The first group of authors observed patients with acute cholecystitis only. Overall conversion rate was about 5% (graph 3)



Graph 2. The conversion rate of individual studies.



Graph 3. Overall conversion rate of the review

We found 12 patient-related risk factors of conversion in laparoscopic cholecystectomy throughout our eligible studies. Three out of them, dense adhesions/fibrosis of GB & Callot's triangle, unclear anatomy at Calot's triangle, and severe forms of inflammation were the most popular causes of conversion in almost all studies. Most of the authors mentioned those three factors with significantly high numbers (tables 2). In three studies, the proportion of converted patients due to the dense adhesions/fibrosis of GB & Callot's triangle made up over 70 percent (see graph 4). 5 out of 22 eligible studies did not mention this factor as a reason for conversion from laparoscopic cholecystectomy. Other causes for conversion were negligible. Mean of abscess for conversion was 6,35% $p=0.260$; Mirizzi syndrome - 3,65%, $p=0.376$; Contracted gallbladder - 2,825%, $p=0.04$; Dilated cystic duct with unsuccessful cholangiogram – 8.85%, $p=0.018$; Thickened gallbladder/grasping inability – 6.95%, $p=0.278$.

Table 2.

Intraoperative patient-related risk factors of conversion from laparoscopic surgery to open surgery.

Authors	Conversion	Patient-related factors of conversion				
		Dense adhesion s/fibrosis of GB & Callot's triangle	Unclear anatomy	Severe inflammation	Biliodigestive fistula	Mirizzi syndrome
El Nakeeb et al	83 (2.5%)	35 (42.2%)	29 (34.9%)		2 (2.4%)	
Gangemi et al	11 (3.87%)	2 (18.2%)	1 (9.1%)	2 (18.2%)		
Licciardello et al	33 (7.9%)	14 (42.4%)		12 (36.3%)	1 (3%)	2 (6%)
Taki-Eldin & Badawy	24 (4.9%)		4 (16.7%)	11 (45.8%)		-
Malik	33 (3.52%)	25 (75.7%)		5 (15.1%)		
Simopoulos et al	94 (5.2%)		70 (74.4%)		6 (6.4%)	1 (1%)

Lo et al	8 (11%)	6 (75%)			1 (12.5%)	
Shamiyeh et al	245 (5.4%)	35 (14.3%)	42 (17.1%)	72 (29.4%)	7 (2.95)	3 (1.2%)
Yajima et al	47 (11.6%)	15 (31.9%)		8 (17%)		-
Ishizaki et al	89 (7.5%)		55 (62%)			4 (4%)
Johanning et al	234 (8.8%)	51 (21.8%)	39 (16.7%)	61 (26.1%)		
Lim et al	56 (27.7%)	20 (35.8%)	26 (46.4%)			
Ercan et al	101 (5%)	58 (57.4%)	20 (19.8%)		3 (3%)	
Kama et al	48 (4.8%)		34 (3.4%)			
Pavlidis et al	98 (7.8%)	37 (37.7%)	1 (1%)	20 (20.4%)	2 (2%)	3 (3%)
Kala et al	82 (1%)	7 (8.5%)		38 (46.3%)	1 (1.2%)	10 (12.2%)
Shamim et al	81 (6.5%)	4 (5%)	44 (54.3%)		1 (1.2%)	
Hadad et al	109 (8.7%)	32 (29.3%)	2 (1.8%)	4 (3.6%)		
Rashid et al	21 (7%)	8 (38.1%)	5 (23.8%)			
Genc et al	163 (3.2%)	118 (76.3%)			4 (2.4%)	1 (0.6%)
Zhang et al	94 (7.4%)	26 (27.75%)	52 (55.3%)			5 (5.3%)
Sikora et al	29 (19.4%)		9 (31%)	9 (31%)		
Total	1783/40 315 (4.6%)	493 (27.6%)	433 (24.3%)	242 (13.5%)	28 (1.5%)	29 (1.6%)
Mean±SD (%)	7,758±5,88 23	37,491±22, 450	29,231±23,0 27	26,29±13,1 895	3,705±3,41 6	4,175±3,75 7
Median (%)	6,75	35,8	21,8	26,1	2,675	3,5
Minimum (%)	1	5	1	3.6	1.2	0.6
Maximum (%)	27.7	76.3	74.4	46.3	12.5	12.2
p value	<0.001*	<0.001*	<0.001*	<0.001*	0,008*	0.016*

*Statistically significant result ($p < 0.05$)

Overall, 11 surgeon-related risk factors of conversion were identified including bile duct injury (4.24%, $p < 0.001$), uncontrollable bleeding (10.06%, $p < 0.001$), duodenal injury (3.41%, $p = 0.06$), colonic injury (0.9%, $p = 0.2$), bile leakage (4.65%, $p = 0.193$), spilled stones of the gallbladder into the abdominal cavity (2.35%, $p = 0.053$), common bile duct exploration due to large CBD stone (3.7%, $p = 0.01$), torn cystic duct (3.4%), failed insertion of trocar or initial laparoscopy failure (4.8%, $p = 0.36$), anesthesia (4.8%) and respiratory disturbances (2.1%) (tables 3).

Table 3. Intraoperative surgeon-related risk factors of conversion from laparoscopic surgery to open surgery

Authors	Conversion	Surgeon-related factors of conversion				
		Bile duct injury	Bleeding	Duodenal injury (perforation)	Common bile duct exploration (due to large CBD stone)	Bile leakage
El Nakeeb et al	83 (2.5%)	2 (2.4%)	6 (7.2%)			
Gangemi et al	11 (3.87%)					

Licciardello et al	33 (7.9%)	2 (6%)	2 (6%)			
Taki-Eldin & Badawy	24 (4.9%)		9 (37.5%)			
Malik	33 (3.52%)	1 (3%)	1 (3%)	3 (9%)		
Simopoulos et al	94 (5.2%)	2 (2.1%)	3 (3.2%)			
Lo et al	8 (11%)					1 (12.5%)
Shamiyeh et al	245 (5.4%)	8 (3.3%)	22 (8.9%)	3 (1.2%)	9 (3.7%)	2 (0.8%)
Yajima et al	47 (11.6%)	5 (10.6%)	8 (17%)	1 (2.1%)		
Ishizaki et al	89 (7.5%)	11 (12%)	15 (17%)		4 (4%)	
Johanning et al	234 (8.8%)		32 (13.7%)			
Lim et al	56 (27.7%)		10 (17%)			
Ercan et al	101 (5%)	9 (8.9%)	10 (9.9%)			
Kama et al	48 (4.8%)	5 (0.5%)	2 (0.2%)	3 (0.3%)	1 (0.1%)	
Pavlidis et al	98 (7.8%)	2 (2%)	10 (10.2%)		4 (4%)	
Kala et al	82 (1%)	5 (6.1%)	6 (7.3%)			
Shamim et al	81 (6.5%)		6 (7.4%)	1 (1.2%)	5 (6.1%)	
Hadad et al	109 (8.7%)	3 (2.7%)	1 (0.9%)			
Rashid et al	21 (7%)	1 (4.7%)	2 (9.5%)	2 (9.5%)		1 (4.7%)
Genc et al	163 (3.2%)	6 (3.7%)	14 (8.5%)	1 (0.6%)	1 (0.6%)	1 (0.6%)
Zhang et al	94 (7.4%)	4 (4.3%)	-		7 (7.4%)	
Sikora et al	29 (19.4%)	1 (3.4%)	4 (13.8%)			
Total	1783/40 315 (4.6%)	119 (6.6%)	163 (9.1%)	14 (0.7%)	31 (1.7%)	5 (0.2%)
Mean±SD (%)	7,758±5,8823	4,246±2,695	10,066±8,394	3,414±4,028	3,7±2,652	4,650±5,563
Median (%)	6,75	3,40	8,70	1,20	4	2,750
Minimum (%)	1	0.5	0.2	0.3	0.1	0.6
Maximum (%)	27.7	10.6	37.5	9.5	7.4	12.5
p value	<0.001*	<0.001*	<0.001*	0,06	0,01*	0,193

*Statistically significant result ($p < 0.05$)

Regarding the equipment-related factors of conversion in laparoscopic surgery, 6 studies had technical issues with a small number of patients when they had to convert the surgery to open cholecystectomy. Other reasons related to this category are rare and insignificant.

DISCUSSION. A total of 22 studies were included in this systematic review. Overall, 40 315 patients were observed in those researches. During this review, we found dense adhesions or fibrosis at Calot's triangle and around GB, unclear anatomy at Calot's triangle, and severe destructive forms of inflammation including empyema, gangrene, and abscess as the most common intraoperative patient-related reasons for conversion from laparoscopic cholecystectomy to open surgery. Furthermore, bile duct injury and uncontrollable bleeding were the most popular intraoperative surgeon-related causes of conversion in laparoscopic cholecystectomy. Equipment-related and other factors of intraoperative conversion from laparoscopy to open surgery were negligible and insignificant according to the results of this review.

There is a significant number of studies in the literature investigating preoperative or predictable risk factors of conversion from laparoscopic cholecystectomy to open surgery. Most of these studies are focused on revealing the predictable reasons for conversion and attempted to offer their prediction models of the conversion risk factors in laparoscopic cholecystectomy. Some authors of the studies offered the Tokyo Guideline 2013 as a perfect prediction model to prevent unwanted conversion (M. Bouassida et al, 2017 M. Yokoe et al, 2013). Another team of authors of the research advocated their version of the prediction model (M.S. Kim et al, 2014).

Yet, there is an obvious lack of papers in the literature that reports on the importance of intraoperative or unpredictable causes of conversion from laparoscopic cholecystectomy to open surgery. This lack is even more serious regarding the secondary studies on this topic. Similar to primary studies, the results of some systematic reviews and meta-analyses have analyzed the papers devoted to the predictable factors of conversion of laparoscopic cholecystectomy. For instance, A.S.Y. Hu et al 2017 focused on body mass index, gall bladder wall thickness, previous history of abdominal surgery, presence of choledocholithiasis, patients' age, patient settings (emergency, elective), blood tests for acute inflammation (alkaline phosphate, white blood cell count, etc.), total bilirubin and other risk factors. This review excluded intraoperative findings and conditions that may lead to conversion to open surgery.

Many studies reported that inflammation, symptomatic gallstones, and fibrosis are significantly more extensive in male genders than in females (S. Yol et al, 2006, A. Gangemi et al, 2017, C. Simopoulos et al, 2005). This is the main reason for the higher conversion rate of laparoscopic cholecystectomy among males. Likewise, in patients with acute gallbladder disease, the rate of conversion

is significantly high due to challenging dissection at thickened and densely adherent structures at Calot's triangle.

This review identified several intraoperative risk factors of conversion that have not been investigated by many secondary studies. These factors include bilio-digestive fistula, spilled gallstones into the abdominal cavity, torn cystic duct, and other uncommon causes of conversion.

Limitations of the review. The review included the papers published in English only. There might be studies reporting intraoperative risk factors of conversion in other languages that were not identified by the authors of this research owing to the language filter. Some studies with very interesting titles and abstracts were not available in full texts. Due to the heterogeneity of included studies in the review, we could not accomplish a meta-analysis.

CONCLUSION. We found a wide range of unpredictable risk factors of conversion in laparoscopic cholecystectomy reported in included studies. Some of those factors are vitally important and very common in surgical practice. Further investigations into the reasons for those risk factors can be most helpful and effective tool for minimization the complications and unwanted conversion in laparoscopic surgery.

REFERENCES

1. "A prospective analysis of 1518 laparoscopic cholecystectomies. The Southern Surgeons Club." (1991). *The New England journal of medicine* vol. 324,16: 1073-8. doi:10.1056/NEJM199104183241601
2. Hu ASY, Menon R, Gunnarsson R, de Costa A. Risk factors for conversion of laparoscopic cholecystectomy to open surgery - A systematic literature review of 30 studies. *Am J Surg.* 2017;214(5):920-930. doi:10.1016/j.amjsurg.2017.07.029
3. Philip Rothman J, Burcharth J, Pommergaard H.C, Viereck S, Rosenberg J. Preoperative Risk Factors for Conversion of Laparoscopic Cholecystectomy to Open Surgery - A Systematic Review and Meta-Analysis of Observational Studies. *Dig Surg.* 2016;33(5):414-423. doi:10.1159/000445505
4. Gangemi A, Danilkowicz R, Bianco F, Masrur M, Giulianotti PC. Risk Factors for Open Conversion in Minimally Invasive Cholecystectomy. *JSLS.* 2017;21(4):e2017.00062. doi:10.4293/JSLS.2017.00062
5. Sutcliffe R.P, Hollyman M, Hodson J, et al. Preoperative risk factors for conversion from laparoscopic to open cholecystectomy: a validated risk score derived from a prospective U.K. database of 8820 patients. *HPB (Oxford).* 2016;18(11):922-928. doi:10.1016/j.hpb.2016.07.015

6. Moher D, Liberati A, Tetzlaff J, Altman D.G; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535. Published 2009 Jul 21. doi:10.1136/bmj.b2535
7. University of York. PROSPERO International Prospective Register of Systematic Reviews. Available at: <http://www.crd.york.ac.uk/PROSPERO/>. (Accessed: May 2019).
8. Kim M.S, Kwon H.J, Park H.W, et al. Preoperative prediction model for conversion of laparoscopic to open cholecystectomy in patient with acute cholecystitis: based on clinical, laboratory, and CT parameters. *J Comput Assist Tomogr*. 2014;38(5):727-732. doi:10.1097/RCT.000000000000116
9. Taki-Eldin A, Badawy A.E. Outcome of laparoscopic cholecystectomy in patients with gallstone disease at a secondary level care hospital. *Arq Bras Cir Dig*. 2018;31(1):e1347. Published 2018 Jun 21. doi:10.1590/0102-672020180001e1347
10. El Nakeeb A, Mahdy Y, Salem A, et al. Open Cholecystectomy Has a Place in the Laparoscopic Era: a Retrospective Cohort Study. *Indian J Surg*. 2017;79(5):437-443. doi:10.1007/s12262-017-1622-2
11. Gangemi A, Danilkowicz R, Bianco F, Masrur M, Giulianotti PC. Risk Factors for Open Conversion in Minimally Invasive Cholecystectomy. *JSLS*. 2017;21(4):e2017.00062. doi:10.4293/JSLS.2017.00062
12. Rashid T, Naheed A, Farooq U, Iqbal M, Barkat N. Conversion of laparoscopic cholecystectomy into open cholecystectomy: an experience in 300 cases. *J Ayub Med Coll Abbottabad*. 2016;28(1):116-119
13. Malik AM. Difficult laparoscopic cholecystectomies. Is conversion a sensible option? *J Pak Med Assoc*. 2015;65(7):698-700
14. Licciardello, A., Arena, M., Nicosia, A., Di Stefano, B., Cali, G., Arena, G., & Minutolo, V. (2014). Preoperative risk factors for conversion from laparoscopic to open cholecystectomy. *European review for medical and pharmacological sciences*, 18(2 Suppl), 60–68.
15. Kala S, Verma S, Dutta G. Difficult situations in laparoscopic cholecystectomy: a multicentric retrospective study. *Surg Laparosc Endosc Percutan Tech*. 2014;24(6):484-487. doi:10.1097/SLE.0b013e31829cebd8
16. Yajima H, Kanai H, Son K, Yoshida K, Yanaga K. Reasons and risk factors for intraoperative conversion from laparoscopic to open cholecystectomy. *Surg Today*. 2014;44(1):80-83. doi:10.1007/s00595-012-0465-5
17. Genc V, Sulaimanov M, Cipe G, et al. What necessitates the conversion to open cholecystectomy? A retrospective analysis of 5164 consecutive laparoscopic operations. *Clinics (Sao Paulo)*. 2011;66(3):417-420. doi:10.1590/s1807-59322011000300009

18. Ercan M, Bostanci EB, Teke Z., Karaman K., Dalgic T, Ulas M, Ozer I, Ozogul YB, Atalay F, Akoglu M. (2010). Predictive factors for conversion to open surgery in patients undergoing elective laparoscopic cholecystectomy. *Journal of laparoendoscopic & advanced surgical techniques. Part A*, 20(5), 427–434. <https://doi.org/10.1089/lap.2009.0457>
19. Shamim M, Memon A.S, Bhutto A.A, Dahri M.M. Reasons of conversion of laparoscopic to open cholecystectomy in a tertiary care institution. *J Pak Med Assoc.* 2009;59(7):456-460.
20. Zhang WJ, Li JM, Wu G.Z, Luo K.L, Dong ZT. Risk factors affecting conversion in patients undergoing laparoscopic cholecystectomy. *ANZ J Surg.* 2008;78(11):973-976. doi:10.1111/j.1445-2197.2008.04714.x
21. Pavlidis TE, Marakis G.N, Ballas K, Symeonidis N, Psarras K, Rafailidis S, Karvounaris D, Sakantamis A.K. (2007). Risk factors influencing conversion of laparoscopic to open cholecystectomy. *J Laparoendosc Adv Surg Tech A.*, 17(4), 414-8. DOI: 10.1089/lap.2006.0178. PubMed PMID: 17705718
22. Lim K.R, Ibrahim S, Tan NC, Lim SH, Tay KH. Risk factors for conversion to open surgery in patients with acute cholecystitis undergoing interval laparoscopic cholecystectomy. *Ann Acad Med Singap.* 2007;36(8):631-635
23. Hadad S.M, Vaidya J.S, Baker L, et al. Delay from symptom onset increases the conversion rate in laparoscopic cholecystectomy for acute cholecystitis [published correction appears in *World J Surg.* 2008 Dec;32(12):2747. Hussain, Kashif [added]]. *World J Surg.* 2007;31(6):1298-1303. doi:10.1007/s00268-007-9050-2
24. Sikora S.S, Kumar A, Saxena R, Kapoor V.K, Kaushik SP. Laparoscopic cholecystectomy--can conversion be predicted?. *World J Surg.* 1995;19(6):858-860. doi:10.1007/BF00299786
25. Ishizaki Y, Miwa K, Yoshimoto J, Sugo H, Kawasaki S. (2006). Conversion of elective laparoscopic to open cholecystectomy between 1993 and 2004. *The British journal of surgery*, 93(8), 987–991. <https://doi.org/10.1002/bjs.5406>.
26. Simopoulos C, Botaitis S, Polychronidis A, Tripsianis G, Karayiannakis AJ. Risk factors for conversion of laparoscopic cholecystectomy to open cholecystectomy. *Surg Endosc.* 2005;19(7):905-909. doi:10.1007/s00464-004-2197-0
27. Kama N.A, Doganay M, Dolapci M, Reis E, Atli M, Kologlu M. Risk factors resulting in conversion of laparoscopic cholecystectomy to open surgery. *Surg Endosc.* 2001;15(9):965-968. doi:10.1007/s00464-001-0008-4

28. Johanning J.M, Gruenberg JC. (1997). "AEIOU: the ABC's" of conversion from laparoscopic to open cholecystectomy. *JSLS: Journal of the Society of Laparoendoscopic Surgeons*, 1(2), 181–183

29. Bouassida M, Chtourou MF, Charrada H, et al. The severity grading of acute cholecystitis following the Tokyo Guidelines is the most powerful predictive factor for conversion from laparoscopic cholecystectomy to open cholecystectomy. *J Visc Surg*. 2017;154(4):239-243. doi:10.1016/j.jviscsurg.2016.11.007

30. Yokoe M, Takada T, Strasberg SM, et al. New diagnostic criteria and severity assessment of acute cholecystitis in revised Tokyo Guidelines. *J Hepatobiliary Pancreat Sci*. 2012;19(5):578-585. doi:10.1007/s00534-012-0548-0

31. Yol S, Kartal A, Vatansev C, Aksoy F, Toy H. Sex as a factor in conversion from laparoscopic cholecystectomy to open surgery. *JSLS*. 2006;10(3):359-363.

Contents

THE EFFECTIVENESS OF THE USE OF PEDAGOGICAL TECHNOLOGIES IN TEACHING UROLOGICAL SUBJECTS Mirzamakhmud A. Shadmanov	5
ANALYSIS OF THE IMMEDIATE RESULTS OF THE MULTIMODAL METHOD OF BREAST CANCER THERAPY IN COMPARISON WITH THE RESULTS OF CONVENTIONAL THERAPY Djakhongir R. Sabirov, Mirza A. Gafur-Akhunov, Oybarchin J. Yusupova	9
THE STUDY OF DIAGNOSTICS AND PREVENTION OF PATHOPHYSIOLOGICAL PARAMETERS AFTER MODERN TREATMENT OF PURULENT-NECROTIC PROCESSES IN DIABETIC FOOT SYNDROME Ulugbek Y.Ergashev, Adkhamjon R. Zokhirov, Khojimurod I. Ernazarov, Ravshankhuja R. Minavarkhyjayev, Nodir M. Malikov	22
COVID-19 ASSOCIATED CAVERNOUS SINUS THROMBOSIS: 2-YEAR FOLLOW-UP EXPERIENCE Erkin N. Bilalov, Okilkhon I. Oripov, Ravshan Z. Umarov, Gulomboy U. Khudaibergenov, Bakhodir E. Bilalov	36
THE STUDY OF PATHOMORPHOLOGICAL DIAGNOSIS OF VITAL ORGANS AFTER MODERN TREATMENT OF DIABETIC FOOT SYNDROME Ulugbek Y.Ergashev, Adkhamjon R. Zokhirov, Khojimurod I. Ernazarov, Ravshankhuja R. Minavarkhyjayev, Behzod A. Abdusalomov	45
INTRAOPERATIVE REASONS FOR CONVERSION OF LAPAROSCOPIC CHOLECYSTECTOMY TO OPEN SURGERY (SYSTEMATIC REVIEW) Khakimov M.Sh., Karimov R.A., Jasmin Sabanovic, Karim Belhaj, Bijendra Patel.....	56
IMPROVING THE TREATMENT OF CORONAVIRUS INFECTION COVID-19 Pulat M. Abilov, Bakhtiyar U. Iriskulov, Ozoda Z. Saydalikhodjaeva, Zukhra N. Boboeva, Sevara B. Azimova, Gulchekhra E. Usmonova.....	69
OPTIMIZATION OF THERAPY FOR GASTROESOPHAGEAL REFLUX DISEASE IN COMORBIDITY WITH FUNCTIONAL DYSPEPSIA Mirvosit M. Karimov, Pulat S. Zufarov, Guzal N. Sobirova, Shahlo S. Aripdjanova, Dildora K. Karimova	77
VITAMINS D AND C IN COVID-19 Dilafroz A. Khatamova, Jakhongir O. Khamdamov	85
MODERN ADVANCES IN THE DIAGNOSIS OF ACUTE CALCULOUS CHOLECYSTITIS (A literature review) Khakimov M.Sh., Karimov R.A., Murodov A.M.	91
USING DIFFERENT TYPES OF ASSESSMENT IN LANGUAGE TEACHING R.O. Ermatova, Z.Kh. Paygamova	105

THE ROLE OF INFLAMMATORY MARKERS AS IL-6, TNF-a, INF-y WITH PSORIATIC ARTHRITIS AND RATES OF COMORBIDITY Khilola T. Mirakhmedova, Gulizebo B. Saidrasulova, Nilyufar R. Mukhsimova.....	110
HEMOGRAM CHANGES IN POSTKOVID SYNDROME IN PREGNANT WOMEN Dilafuz L. Zaynutdinova, Shaira A. Babadjanova	118
PREDICTION OF LONG-TERM NEUROLOGICAL CONSEQUENCES OF CORONAVIRAL INFECTION USING NEUROTROPIC AUTOANTIBODIES Firuza Kh. Inoyatova, Gulnora K. Rakhmatullaeva, Nigina A. Vakhobova, Umida S. Salikhodjaeva	124
ANALYSIS OF THE PROCESS OF DEEP MACHINE LEARNING BASED ON THE RESULTS OBTAINED FOR PRIMARY DIAGNOSTICS OF GASTROENTEROLOGICAL DISEASES Rustam Yaxshiboyev, Bahodir Muminov, Erkin Ermetov, Abdunazar Jurayev.....	138
EFFICACY OF ENDOSURGICAL CORRECTION OF FEMALE OVARIAN INFERTILITY G.A. Tanish, G.S. Babadjanova.....	148
COMPLEX SURGICAL TREATMENT OF PATIENTS WITH HIGH-GRADE BRAIN TUMORS Sh.N. Tashmatov, R.T. Kadyrbekov, G.A. Alikhodjaeva, T.M. Akhmediev, Z.Sh. Shamuratov	153
ASPECTS OF TEACHING STUDENTS TELEMEDICINE SKILLS IN MEDICAL UNIVERSITIES Bobir T. Khalmukhamedov, Nargiza M. Nurillaeva	163
THE FREQUENCY AND SEVERITY OF COMPLICATED PYELONEPHRITIS AFTER ENDOSCOPIC INTERVENTIONS FOR UROLITHIASIS AND ITS RISK FACTORS F.A. Akilov, Sh.I. Giyasov, A.R. Ruzibaev, I.B. Ziyaev, A.A. Rakhimbaev	172
PROBLEMS WITH REDUCED BONE DENSITY IN SYSTEMIC SCLERODERMA Sevara M. Mukhammadieva, Alibek A. Khudoynazarov.....	186



MUHARRIRIYAT VA NASHRIYOT BO'LIMI

Volume – 13,62 usl. printer. Circulation – 100. Format 60x84. 1/8.
Listening means «Times New Roman». Printed in TMA editorial and publisher department.
100109. St. Farabi 2, Tel.: (998 71)214-90-64, e-mail: rio-tma@mail.ru