

O'ZBEKISTON RESPUBLIKASI
OLIIY TA'LIM, FAN VA INNOVATSIYALAR VAZIRLIGI
FARG'ONA DAVLAT UNIVERSITETI

**FarDU.
ILMIY
XABARLAR**

1995 yildan nashr etiladi
Yilda 6 marta chiqadi

1-2023

**НАУЧНЫЙ
ВЕСТНИК.
ФерГУ**

Издаётся с 1995 года
Выходит 6 раз в год

Muassis: Farg'ona davlat universiteti.

“FarDU. Ilmiy xabarlar – Научный вестник. ФерГУ – Scientific journal of the Fergana State University” jurnali bir yilda olti marta elektron shaklda nashr etiladi.

Jurnal pedagogika, filologiya, tarix, falsafa, siyosat, kimyo, biologiya hamda geografiya fanlari bo'yicha O'zbekiston Respublikasi Oliy attestatsiya komissiyasining doktorlik dissertatsiyalari asosiy ilmiy natijalarini chop etish tavsiya etilgan ilmiy nashrlar ro'yxatiga kiritilgan.

Jurnaldan maqola ko'chirib bosilganda, manba ko'rsatilishi shart.

O'zbekiston Respublikasi Prezidenti Administratsiyasi huzuridagi Axborot va ommaviy kommunikatsiyalar agentligi tomonidan 2020-yil 2-sentabrda 1109 raqami bilan ro'yxatga olingan.

Muqova dizayni va original maket FarDU tahririy-nashriyot bo'limida tayyorlandi.

Tahrir hay'ati

Bosh muharrir
Mas'ul muharrir

SHERMUHAMMADOV B.SH.
ZOKIROV I.I.

FARMONOV Sh. (O'zbekiston)	JEHAN SHAHZADAH NAYYAR (Yaponiya)	G'ULOMOV S.S. (O'zbekiston)
BEZGULOVA O.S. (Rossiya)	LEEDONG WOOK (Janubiy Koreya)	BERDISHEV A.S. (Qozog'iston)
RASHIDOVA S. (O'zbekiston)	A'ZAMOV A. (O'zbekiston)	KARIMOV N.F. (O'zbekiston)
VALI SAVASH YYELEK (Turkiya)	KLAUS XAYNSGEN (Germaniya)	CHESTMIR SHTUKA (Slovakiya)
ZAYNOBIDDINOV S. (O'zbekiston)	BAXODIRXONOV K. (O'zbekiston)	TOJIBOYEV K. (O'zbekiston)
AXTARIYEVA R. (Rossiya)	RAKETIN O. (Rossiya)	TURAYEV A. (O'zbekiston)

Tahririyat kengashi

IJTIMOYIY GUMANITAR FANLARI BO'YICHA		ANIQ VA TABIIY FANLAR BO'YICHA	
EGAMBERDIYEVA T. (O'zbekiston)	FarDU, p.f.d.	MAMAJONOV M. (O'zbekiston)	FarDU, f.f.b.f.d.(PhD)
XONKELDIYEV Sh. (O'zbekiston)	FarDU, p.f.d.prof.	MAMAJONOV A. (O'zbekiston)	FarDU, f.f.b.f.d (PhD)
TOJIBOYEVA H. (O'zbekiston)	Ped.ilm.tad.instituti. p.f.d.(Dsc)	ISOMIDDINOV F. (O'zbekiston)	FarDU, f.f.n.
O'RINOVA N. (O'zbekiston)	FarDU, p.f.d.prof.	SABIRDINOV A. (O'zbekiston)	FarDU, f.f.d.prof.
USMONOV B. (O'zbekiston)	FarDU, p.f.d.	RAHIMOV Z. (O'zbekiston)	FarDU, f.f.d.
DAVLYATOVA G. (O'zbekiston)	FarDU, p.f.n.	XOSHIMOVA N. (O'zbekiston)	FarDU, f.f.b.f.d(PhD)
G'OFUROV A. (O'zbekiston)	FarDU, i.f.d.prof.	MIRZAYEVA D. (O'zbekiston)	FarDU, f.f.b.f.d(PhD)
XONKELDIYEVA G. (O'zbekiston)	FarDU, i.f.d.	ANIQ VA TABIIY FANLAR BO'YICHA	
HOLMATOV B. (O'zbekiston)	FarDU, I.f.n.	O'RINOV A.Q. (O'zbekiston)	FarDU, f.m.f.d.prof.
XAKIMOV N. (O'zbekiston)	FarDU, f.f.d. dots.	KARIMOV Sh.T (O'zbekiston)	FarDU, f.m.f.d.dots.
ISOMIDDINOV M. (O'zbekiston)	FarDU, t.f.d.	OTAJONOV S. (O'zbekiston)	FarDU, f.m.f.d.prof.
USMONOV B. (O'zbekiston)	FarDU, t.f.d (DSc)	RASULOVA R. (O'zbekiston)	FarDU, f.m.f.d.prof.
MAXMUDOV O. (O'zbekiston)	FarDU, t.f.d.prof (PhD)	ONARQULOVA K. (O'zbekiston)	FarDU, f.m.f.d.prof.
ALIMOVA N. (O'zbekiston)	FarDU, t.f.d(PhD)	SHUHRATOV SH. (O'zbekiston)	FarDU, t.f.b.f.d (PhD)
JO'RAYEV H. (O'zbekiston)	FarDU, t.f.d(PhD).dots.	ASQAROV I. (O'zbekiston)	AnDU, k.f.d.prof.
MAMATOV M. (O'zbekiston)	FarDU, t.f.d(PhD).dots.	IBRAGIMOV A.(O'zbekiston)	FarDU, k.f.d.prof.
AHRORQULOVA A. (O'zbekiston)	FarDU, t.f.b.f.d(PhD)	MAMATQULOVA S. (O'zbekiston)	FarDU, k. f.b.f.d (PhD)
G'ANIYEV B. (O'zbekiston)	FarDU, f.f.d.prof.	ABDUG'ANIYEV O. (O'zbekiston)	FarDU, g.f.n.
MAMAJONOV A. (O'zbekiston)	FarDU, f.f.d.prof.	AXMADALIYEV Yu. (O'zbekiston)	FarDU, g.f.d.prof.
ISKANDAROVA Sh. (O'zbekiston)	FarDU, f.f.d.prof.	YULDASHEV G. (O'zbekiston)	FarDU, q.x.f.d.
SHUKUROV R. (O'zbekiston)	FarDU, f.f.d.prof.	ISAG'ALIYEV M. (O'zbekiston)	FarDU, b.f.d. dots.
YULDASHEVA D. (O'zbekiston)	FarDU, f.f.doktori.	SHERALIYEV B. (O'zbekiston)	FarDU, b.f.d.(PhD)
UMAROVA N. (O'zbekiston)	FarDU, f.f.doktori.	SHERMATOV M. (O'zbekiston)	FarDU, b.f.n.

Muharrir: Sheraliyeva J.

Tahririyat manzili:

150100, Farg'ona shahri, Murabbiylar ko'chasi, 19-uy.
Tel.: (0373) 244-44-57. Mobil tel.: (+99891) 670-74-60
Sayt: www.fdu.uz. Jurnal sayti

Bosishga ruxsat etildi:

Qog'oz bichimi: - 60x84 1/8

Bosma tabog'i:

Ofset bosma: Ofset qog'ozi.

Adadi: 10 nusxa

Buyurtma №

FarDU nusxa ko'paytirish bo'limida chop etildi.

Manzil: 150100, Farg'ona sh., Murabbiylar ko'chasi, 19-uy.

Farg'ona, 2023

I.R.Asqarov, M.M.Mo'minov, U.Sh.Xusanov

Gulxayri(<i>althaea officinals</i> l) o'simligini kimyoviy tarkibi, gulxayri moyini elementlar analizi va uning xalq tabobatidagi axamyati	120
---	-----

BIOLOGIYA**Sh.X.Yusupova, I.I.Zokirov**

No'xat agrotsenozi zararli entomofaunasining ekologo-faunistik tahlili (shimoliy Farg'ona misolida)	127
---	-----

K.Zokirov, A.K.Xusanov, O.T.Sobirov, M.F.Xafizddinov, D.A.Saidjaxonova, S.T.Tillayeva, A.A.Kozimov

Sharqiy Farg'ona sharoitida terak qabariq qalqondori (<i>diaspidiotus slavonicus</i> (green, 1934)ning biologik va zoogeografik xususiyatlariga oid	135
--	-----

Z.J.Isomiddinov, D.A.Mirzaliyeva

Xushbo'y shivit (<i>anethum graveolens</i> l.) O'simligining biokimyoviy xossalari	143
---	-----

F.I.Xalmetova, X.S.Axmedov, S.N.Buranova, A.N.Botirbekov

Reaktiv artriting genetik jihatlari	146
---	-----

M.R.Shermatov

Farg'ona vodiysi agroekotizimlari tangachaqanotli hasharotlarining (insecta: lepidoptera) zoogeografik tahlili	150
--	-----

K.Z.Yakhyeva, F.F.Xoltayeva, K.K.Aliyeva

Chaqaloqlarda buyrak patologiyasi sabalari	157
--	-----

M.A.Raximov

Mollarni go'shtga boqishda genetik imkoniyatlaridan samarali foydalanish	161
--	-----

M.X.Mirraximova, N.Y.Nishonboeva

Genining polimorfizmi atopik dermatitda	165
---	-----

G.M.Zokirova

Farg'ona vodiysi sharoitida <i>cinara tujafilina</i> (del guercio, 1909) shirasining (homoptera, lachnidae) biologiyasi	169
---	-----

E.A.Botirov

Janubiy Farg'onaning <i>agrotis</i> avlodi tunlam kapalaklari faunasi va ekologik xususiyatlari	173
---	-----

G.M.Duschanova, N.A.Sobirova, D.A.Abdullayev

Toshkent botanika bog'i sharoitida <i>Eremurus lactiflorus</i> O. Fedtsch. (<i>Xanthorrhoeaceae</i>) o'simligi bargining strukturaviy xususiyatlari	179
---	-----

F.M.Abdualiyeva, Sh.S.Xushmatov

Andijon shahrida joylashgan №1-son maktab o'quvchilarining (1-11 sinf) anfimova testi asosida intellektual rivojlanish darajasi tahlili	185
---	-----

GEOGRAFIYA**Y.I.Ahmadaliyev, X.A.Abdualiyev**

Aholining hududiy taqsimlanishini tahlil qilishida zamonaviy iqtisodiy-geografik tadqiqotlarning zarurati	191
---	-----

ILMIY AXBOROT**G.M.Mansurov**

Nemis tili darslarida til o'yinlari yordamida suhbatlashish qobiliyatlarini rivojlantirishni o'rganish	196
--	-----

N.A.Sharopova

Qashqadaryo viloyati umumta'lim muassasalari moddiy-texnika bazasini mustahkamlash tadbirlari va ularning natijasi	200
--	-----

O.A.Maniyozov, A.A.Bozorqulov, O.S.Isomiddinova

Ta'lim jarayonida birinchi tartibli chiziqli oddiy differensial tenglamalarni yechimini maple dasturida topish	203
--	-----

D.O'.Qarshiyeva

Ona tili va adabiyot o'qituvchilarining kvest texnologiyasi asosida darslarni tashkil etish kompetentligini rivojlantirish	207
--	-----

G.B.Nafasova, B.S.Abdullayeva

Bo'lajak fizika o'qituvchilarining ilmiy-mantiqiy dunyoqarashini shakllantirish	212
---	-----

U.I.Nosirova, L.A.Ernazarova

Nasriy matnlardagi sintaktik-stilistik figuralarning pragmatik jihatlari	216
--	-----

O.I.Yadgarova

Shaxsni o'rganishda proyektiv metodikalarning afzalliklari tahlili	220
--	-----

UDK: 616.085-72-002.77: 616-002.77

REAKTIV ARTRITNING GENETIK JIHLTLARI

GENETIC ASPECTS OF REACTIVE ARTHRITIS

ГЕНЕТИЧЕСКИЕ АСПЕКТЫ РЕАКТИВНОГО АРТРИТА

**Khalmetova Feruza Iskandarovna¹, Axmedov Xalmurad Sadullayevich²
Buranova Sagdiyana Nasurullayevna³, Botirbekov Aburashid Nazarkulovich⁴**

¹**Khalmetova Feruza Iskandarovna** – Tashkent Medical Academy, senior lecturer (PhD)
²**Axmedov Xalmurad Sadullayevich** – Tashkent Medical Academy, professor
Buranova Sagdiyana Nasurullayevna – Tashkent Medical Academy, assistant (PhD)
Botirbekov Aburashid Nazarkulovich – Tashkent Medical Academy, assistant

Annotatsiya

Reaktiv artrit - bo'g'imlarning eng keng tarqalgan surunkali yallig'lanish kasalliklaridan biri. Reaktiv artrit bilan og'riqan bemorlar revmatologik shifoxonalarda bemorlarning taxminan 10% ni tashkil qiladi va kasallikning surunkali shaklida bo'g'imlarning funksional qobiliyatining sezilarli darajada yo'qolishi va og'ir asoratlar paydo bo'lishi mumkin, bu esa bemorlarning nogironligiga olib keladi. Reaktiv artritga bo'lgan qiziqish, shuningdek, bir qator bemorlarda, ayniqsa, jarayon surunkali bo'lsa, bo'g'imlarda ankilozlanishgacha bo'lgan destruktiv jarayonlarni rivojlanishi bilan bog'liq. Reaktiv artritga chalingan bemorlarning bo'g'imlarida va umurtqa pog'onasida destruksiyaning rivojlanishi o'z navbatida bemorlarning hayot sifatiga salbiy ta'sir etishi, ayniqsa navqiron yoshdagi insonlarning xastalanishi, ularning ish qobiliyatining pasayishi va nogironlik ko'rsatkichining ortib borishi hanuzgacha dolzarb muammo sifatida namoyon bo'lmoqda.

Аннотация

Среди хронических воспалительных заболеваний суставов одним из наиболее часто встречающихся является реактивный артрит. Пациенты реактивным артритом составляют около 10% больных ревматологических стационаров, причем при хронической форме заболевания может развиваться значительная утрата функциональных способностей суставов и возникновение тяжелых осложнений, что приводит к инвалидизации пациентов. Интерес к реактивному артриту связан также с тем обстоятельством, что у ряда больных, особенно при хронизации процесса развиваются деструктивные процессы в суставах вплоть до анкилозирования. Развитие деструкции суставов и позвоночника у пациентов с реактивным артритом, в свою очередь, отрицательно сказывается на качестве жизни пациентов, особенно у молодых людей, а также снижение их трудоспособности и увеличение заболеваемости остается актуальной проблемой.

Abstract

Reactive arthritis is one of the most common chronic inflammatory diseases of the joints. Patients with reactive arthritis make up about 10% of patients in rheumatological hospitals, and in the chronic form of the disease, a significant loss of the functional abilities of the joints and the occurrence of severe complications can develop, which leads to disability of patients. Interest in reactive arthritis is also associated with the fact that a number of patients, especially when the process is chronic, develop destructive processes in the joints up to ankylosing. The development of destruction of the joints and spine in patients with reactive arthritis, in turn, adversely affects the quality of life of patients, especially in young people, as well as a decrease in their ability to work and an increase in morbidity remain an urgent problem.

Kalit so'zlar: reaktiv artrit, citokinlar, spondiloartrit, bo'g'im sindromi, IL-17A rs2275913, IL 18 gen.

Ключевые слова: реактивный артрит, цитокины, спондилоартрит, суставной синдром, ИЛ-17A rs2275913, ген ИЛ 18.

Key words reactive arthritis, cytokines, spondyloarthritis, joints syndrome, IL-17A rs2275913, IL-18 gene

INTRODUCTION. Reactive arthritis (ReA) is a "sterile" inflammatory joint disease in which infectious agents and their antigens are absent in the synovial fluid and synovial membrane of patients. ReA is a seronegative spondyloarthritis, which meets the criteria of the European Group for the Study of Spondyloarthritis. With the introduction into clinical practice in the early 90s of the reverse transcriptase polymerase chain reaction (RT-PCR) method, small amounts of Ch.trachomatis DNA and RNA were found in the joint cavity. The presence of these nucleic acids, which have a very short lifetime in tissues (several minutes), indicated the possibility of transcription and, consequently, active reproduction of bacteria in the joint cavity.

Later it was discovered that the joint is not normally a sterile environment. Thus, the DNA of various microorganisms was found in biopsies of the synovial membrane of healthy people and patients with osteoarthritis (in 9 and 20%, respectively). Genetic factors in the pathogenesis of ReA are being actively studied. In particular, a close association of the disease with the HLA-B27 antigen has been proven. There are several points of view about its role in the development of ReA.

MATERIALS AND METHODS. According to one of them, the HLA-B27 antigen is an antigen-presenting molecule capable of presenting arthritic peptides to cytotoxic CD8 T-lymphocytes. The object of lysis in this case is probably chondrocytes localized both in the cartilage tissue and in the entheses, where the inflammatory process also develops. According to Y. Sobao et al. (1999), for some bacteria (*Yersinia*, *Salmonella*), the presence of HLA-B27 is a factor that significantly facilitates their invasion into synovial membrane cells. In 2000, R.A. Colbert et al. a new theory of the pathogenesis of ReA was proposed, the essence of which is that under the influence of lipopolysaccharides of the cell wall of causal microorganisms during antigenic processing in human monocytes, an incorrect "assembly" of heavy chains of HLA-B27 is possible, which leads to the accumulation of HLA-B27 molecules in the endoplasmic network of these cells.

At the same time, despite the large number of studies conducted in genomic medicine devoted to the study of the mechanisms of development of reactive arthritis, the final answer to this question has not yet been formulated.

Endogenous immunomodulators, that is, cytokines are important in the regulation of almost all body systems. Compared with exogenous modulators (chemical, bacterial or plant origin), cytokines exert their abilities through specific receptors and manifest themselves as natural regulators of the functional activity of various types of tissue. This manifests itself due to the prospects for the use of cytokines in the treatment and prevention of various immunopathological diseases, as well as for the generation of various types of cells during cellular immunotherapy.

RESULTS. It is known that T-helper 17 (Th17) plays an important role in autoimmune processes, destruction of joints and connective tissue. Interleukin-17A (IL-17A) is one of six members of the IL-17 family, which includes IL-17A-F, which are cytokines produced by Th17 cells and other related immune cells. IL-17A stimulates the production of many factors, such as tumor necrosis factor-alpha (TNF- α), IL-6 and IL-1 β , which play an important role in the inflammatory response.

It was studied by Shen L. that the relationship between six IL-17A polymorphisms and the risk of arthritis is the most suitable variant – IL-17A rs2275913, G-197A. However, a meta-analytical study conducted by Chen P. The results showed a significant relationship between IL-17A rs2275913 polymorphism and susceptibility to arthritis in the general population (allele model A versus G; heterozygous model GA versus GG); homozygous model AA versus GG; dominant model GA+ AA versus GG). When analyzing subgroups, IL-17A rs2275913 polymorphism was significantly associated with the risk of arthritis in Europeans (allele model A versus G; heterozygous model GA versus GG; dominant model GA + AA versus GG), but not in Africans or Americans. Since the results of these studies still remain inconclusive. And also according to the study of Marco A. Rocha Loures detected TNF and IL17 genotypes independently of these factors, which may provide more convincing evidence of a link between these polymorphisms and susceptibility to spondyloarthritis, AS disease and psoriatic arthritis. Moreover, the incidence is relatively severe in patients with spondyloarthritis bearing the IL17A A/G genotype and the A allele, as well as the IL17F T/C genotype and the C allele. Many people know that the association of these genetic polymorphisms with spondyloarthritis has not yet been proven.

As a result of many studies conducted, it was found that IL 18 has an important role in the humoral immune response, thereby its significantly higher serum level in patients with Spondyloarthritis. IL 18 belongs to the IL1 family, because of the homology with the IL1 family, IL18 was named IL1u, but as

it turned out later, IL18 does not bind to the IL 1 type I receptor and is therefore isolated as a new cytokine, IL18.

According to Yakushenko E.V.'s research, IL 18 is produced mainly by macrophages, including Kupffer liver cells, and dendritic cells. The presence of IL-18 mRNA is detected in many

cell types, in particular in CD4(+), CD8(+), B- and NK cells. In addition, IL-18 mRNA in humans is found in skeletal muscle cells, keratinocytes, myelomonocytic hematopoietic and human cell lines. It has been proved that in the MyD88-mediated signal activating the inflammatory response cascade, the effects of IL-18 are realized through interaction with a receptor consisting of an auxiliary protein of the IL-18 receptor (IL-18RAP) and the IL-18 receptor protein (IL-18R1). The IL-18RAP/IL-18R1 receptor complex forms a signaling chain and mediates signal transduction initiated by IL-18. Polymorphism of IL18RAP and IL-18R1 genes is associated with a variety of immuno-mediated diseases, which suggests the extreme importance of the balance of IL-18 signaling pathways in connection with the activation of innate and acquired immune response. The expression of the IL 18 gene is so widely represented in various cell types, indicating the participation of this cytokine not only in the formation of the immune response, but also in the regulation of other physiological processes in various tissues and organs. IL 18 has various effects on immunocompetent cells. The main effect of IL 18 is the induction of IFN γ T- and NK production by cells, which is why IL 18 was originally called the IFN γ inducing factor.

The analysis of the amino acid sequence showed the similarity of the structure of IL18 and IL1 β by 18%, IL18 and IL1a or IL1PA by about 15%. Both cytokines have a unique β -folded structure. IL1 β and IL18 are synthesized initially as precursor molecules, then a mature biologically active peptide is produced using the IL1 β converting enzyme (caspase 1). However, despite the partial similarity of the structure of IL1 and IL18, the specific biological functions and receptor systems of these two cytokines are different. If IL18 activates the maturation of type 1 T helper cells (Th1) and does not affect the proliferation of type 2 T helper cells (Th2), whereas IL1 does not affect the proliferative response and production of type 1 cytokines and induces the proliferation of type 18 Th2, its receptor and IL 18 binding protein are a polymorphic structure that is formed as at the level of allelic polymorphism of the genes of these proteins. According to the literature, IL 18 genes were found in more than 9 allelic variants, as well as in IL 18 binding protein more than 11 allelic variants. The presence of any allelic variant of IL 18 or IL 18 binding protein is associated with an increase in the frequency of various pathological processes.

It is known according to K.F.Nolan, the gene encoding IL 18 is located on chromosome 11, locus 1q22.2-q22.3., has 6 exons, 5 introns and one promoter. Despite the fact that the exons and intron-exon boundaries of IL18 were sequenced repeatedly in different populations, no non-synonymous SNPs were found in the gene structure. In addition, no SNPs were found that could interfere with mRNA splicing. This indicates a high conservatism of the gene, most likely directly related to the regulation of the inflammatory response. However, there is a variation in the gene structure within the 5'-untranslated region (UTR) and 3'UTR, which can cause differences in the translation rate and stability of mRNA, as well as changes in the proximal promoter, which affects the transcription rate.

DISCUSSION. However, in order to further determine the role of polymorphic variants and the significance of variations in serum concentrations in the pathogenesis of the progression of diseases of the spondyloarthritis groups, it is necessary to prospectively monitor patients with reactive arthritis, as well as individuals who carry risky genotypes, but do not have joint syndrome at the time of examination.

REFERENCES

1. Аннунциато Ф., Косми Л., Лиотта Ф. и др. Т хелперные клетки типа 17 происхождение, особенности и возможная роль при ревматических заболеваниях.: *Nat Rev Rheumatol*, 2009; 5. – стр. 325–331.
2. Корн Т., Беттелли Э., Оукка М. и др. IL-17 и клетки Th17. *Annu Rev Immunol*, 2009; 27. -Стр. 485–517.
3. Лиля А.М., Гапонова Т.В. Реактивные артриты: особенности патогенеза и терапевтическая тактика. *РМЖ*, 2010;27. – стр. 1663.
4. Понасенко А.В., Цепочкина А.В., Хуторная М.В., Долгих В.А., Малышев И.Ю., Барбараш О.Л. Полиморфизм генов ИЛ 18 гаp и ИЛ 18 r1 ассоциирован с рисками развития инфаркта миокарда у пациентов со стабильной формой ишемической болезни сердца. *Трансляционная медицина*, 2018.Том 5 №4. – стр. 125–131.
5. Шэнь Л., Чжан Х., Ян Т. и др. Связь между полиморфизмом интерлейкина 17A и предрасположенностью к ревматоидному артриту у населения Китая. *Джин* 2015; 566 18–22.
6. Якушенко Е.В. Интерлейкин-18: биологические эффекты и перспективы клинического применения: автореф. дис. д-ра мед. наук 14.03.09 / Якушенко Е.В. – Новосибирск, 2012.
7. Якушенко Е.В., Лопатникова Ю.А., Сенников С.В. Интерлейкин-18 и его роль в иммунном ответе. *Медицинская Иммунология* 2005, Т.7, No 4, стр 355-364.

BIOLOGIYA

8. Abduraxmanova N. M. B., Axmedov X. S. Reactive arthritis-a modern view of the problem //Журнал биомедицины и практики. – 2021. – Т. 6. – №. 1.
9. Abdurakhmanova N. M. et al. MODERN METHODS OF TREATMENT OF PATIENTS WITH ANKYLOSING SPONDYLITIS //International Journal of Advance Scientific Research. – 2022. – Т. 2. – №. 11. – С. 112-118.
10. Chen P, Li Y, Li L, Zhang G, Zhang F, Tang Y, Zhou L, Yang Y, Li J. Association between the interleukin (IL)-17A rs2275913 polymorphism and rheumatoid arthritis susceptibility: a meta-analysis and trial sequential analysis. *J Int Med Res.* 2021 Oct; 49(10):3000605211053233. doi: 10.1177/03000605211053233. PMID: 34704484; PMCID: PMC8554571.
11. Cohen Susanne E., Bodmer Hellen C. Cytotoxic T lymphocytes recognize and lyse chondrocytes under inflammatory, but not non-inflammatory conditions. *Immunology* 2003; 109:8–14.
12. Colbert R.A. HLA-B27 misfolding: a solution to the spondyloarthropathy conundrum? *Mol Med Today* 2000;6:224–230.
13. Hammer M., Nettelbreker E., Hopf S., Schmitz E., Porshke K., Zeidler H. Chlamidial rRNA in the joints of patients with Chlamydia-induced arthritis and undifferentiated arthritis. *Clin Exp Rheumatol* 1992;10:63–66.
14. J. R. Vidal-Castiñeira, A. López-Vázquez, R. Diaz-Peña et al., "A single nucleotide polymorphism in the *IL17ra* promoter is associated with functional severity of ankylosing spondylitis," *PLoS One*, vol. 11, no. 7, article e0158905, 2016.
15. Khalmetova F. I., Akhmedov K.S., Razakova F.S. Comparative Analysis of the Clinical Presentation of Reactive Arthritis // *American Journal of Medicine and Medical Sciences.* – 2021. –Vol. 11(1). 75-78.
16. Khalmetova F. I., Akhmedov K.S., Buranova S.N. [The role of imaging techniques in the assessment of structural changes in the joint in reactive arthritis](#) - Academia Globe: Inderscience Research, 2022
17. Marco A. Rocha Loures, Luciana C. Macedo, Denise M. Reis, Camila F. Oliveira, Jean L. Meneguetti, Gabriela F. Martines, Janisleya S. F. Neves, Eliana de Souza, Ana M. Sell, Jeane E. L. Visentainer, "Influence of TNF and IL17 Gene Polymorphisms on the Spondyloarthritis Immunopathogenesis, Regardless of HLA-B27, in a Brazilian Population", *Mediators of Inflammation*, vol. 2018, Article ID 1395823, 7 pages, 2018.
18. Schumacher H.R. Jr, Arayssi T., Crane M., et al. Chlamydia trachomatis nucleic acids can be found in the synovium of some asymptomatic subjects. *Arthritis Rheum* 1999;42:1281–1284.
19. Sibilia J., Libmach F-X. Reactive arthritis or chronic infectious arthritis? *Ann Rheum Dis* 2002;61:580–587.
20. Sobao Y., Tsuchiya N., Takiguchi M., et al. Overlapping peptide-binding specificities of HLA-B27 and B39. *Arthritis Rheum* 1999;42:175–181.
21. Mirrakhimova M.K. Improving methods of treatment of atopic pathology in children // *Journal of Critical Reviews.* -№12. -p. 190-192
22. Mirrakhimova M.H., Nishanbaeva N.Y., Clinical Manifestations Of Connective Tissue Dysplasia In Children With Glomerulonephritis // *Journal of Pharmaceutical Negative Results/Volume 13/Special Issue 9 | 2022,rr.4203-4205.*
23. Nishanbayeva N.Yu., Mirrakhimova M.X. Bolalarda atopik dermatitda oshqozon ichak traktidagi klinik laborator o'zgarishlarni aniqlash, tashxislash va davolash tamoyillarini takomillashtirish //«Tibbiyotda yangi kun» 6(38/1)2021 ISSN 2181-712X. EISSN 2181-2187 pp.720-726.