

АСТАНА МЕДИЦИНАЛЫҚ ЖУРНАЛЫ



Специальный выпуск
Том 116, 2023



Астана - 2023

ISSN (print) 1562-2940
ISSN (online) 2799-1293

Министерство здравоохранения
Республики Казахстан



*Астана
медициналық
журналы*

*Astana Medical
Journal*

*Специальный выпуск,
Том 116, 2023*

*Ежеквартальный
научно-практический журнал
Собственник:
НАО "Медицинский университет
Астана"*

ГЛАВНЫЙ РЕДАКТОР
Надыра К.Т.

ЗАМ. ГЛАВНОГО РЕДАКТОРА
Кобко В.В., д.м.н.

ИСПОЛНИТЕЛЬНЫЙ РЕДАКТОР
Цой О.Г., д.м.н., профессор

ИСПОЛНИТЕЛЬНЫЙ РЕДАКТОР
Оразова Е.Ү., PhD

РЕДАКЦИОННАЯ КОЛЛЕГИЯ

Тайгулов Е.А., д.м.н., профессор
(Астана, Казахстан)

Кожахметов С.К., к.м.н.,
ассоциированный профессор
(Астана, Казахстан)

Рустемова К.Р., д.м.н., профессор
(Астана, Казахстан)

Жусупова Г.К., к.м.н.,
(Астана, Казахстан)

Сабитова А.Н., Бактиева Е.Е., Кефаловицкая Е.В. РЕЗУЛЬТАТЫ ПРИМЕНЕНИЯ КРИТЕРИЕВЫХ ШКАЛА ОЦЕНКИ ПРИ ОСТРЫХ МАКСИМ СЧЕДУРАЛЬНЫХ ГЕМАТОМ 11
Золотницкий В.В., Борисовская О.С., Громовикова Г.А., Думчук Т.В., Панова Е.Г., Хомичевская И.С. КЛИНИЧЕСКИЙ СЛУЧАЙ ЗОЛОТЦОВОГО ВНУТРЕННЕГО ПОДДОЛГОВОЙ АРТЕРИИ С ПРИМЕНЕНИЕМ МЕДИКО-ЭМОБЛЮКУЦИОННОЙ СИСТЕМЫ 12
Рыбаковская И.А., Абдуллаева Е.Е., Абдуллаев Е.Е., Абдуллаев Е.Е., Байдукова Е.Е. МАЛОИНВАЗИВНЫЙ СТЕРЕОТАКСИЧЕСКИЙ МЕТОД ЛЕЧЕНИЯ ГЕМОРРАГИЧЕСКОГО ИНСУЛЬТА: КЛИНИЧЕСКИЙ СЛУЧАЙ 13
Каримова А.Е., Чистякова А.Е., Мухаметзянова Е.А. ДИАГНОСТИКА И ЛЕЧЕНИЕ ОДИНАКОВО-ФИБРОЗНОЙ ОБУХОВОЙ ВЛЕМПЫ 14
Широкова Н.Н., Гильзина Г.Н., Бактиева Е.Е., Сабитова А.Н., Жумабекова Е.Е., Жаркин Д.М. ПЕРВЫЕ КЛЫНЫЕ ИСХОДЫ У ВЪЗРЕСШИХ С ВОРОНОВИЧЕВСКОЙ ИНФИКЦИЕЙ 15
Богданов В.Т., Тодорян Г.А., Громовикова Г.С., Демидовская Е.С. УДОБНОСТИСВОИСТАВЛЕННЫЙ ГЕЛЬ НА ОСНОВЕ РЕКОМЕНДАЦИИ АНТИГЕНЕВА ЧЕЛОВЕКА КАК СПОСОБ УСИЛЕНИЯ ПРОТИМЕНОВИРОВОГО, ПРОТИВОВОСПАТИЛЬНОГО И РАНОЛИДЕЛЮВИДНОГО ДЕЙСТВИЙ 16
Богданов В.Т., Жиганова Е.Х., Кузьмин А.В. БЫТ ПРИМЕНЕНИЯ CASE-BASED LEARNING (CBL) НА ЗАНЯТИЯХ ПО ДИСЦИПЛИНЕ «ОФТОЛОГИЯ» 17
Кулакова Е.М., Талгасова Е.А., Ибрагимова Х.Х., Альберт А.А. ИНТЕГРИРОВАННАЯ ПРОГРАММА ОБУЧЕНИЯ МАСТЕРСКОМ С ИСПОЛЬЗОВАНИЕМ СИМУЛЯЦИОННЫХ ТЕХНОЛОГИЙ 18

CONTENT

Reviews

Джарипанов З.А., Ахмедзода Р., Бондаренко Е. MODERN VISION OF THE TREATMENT OF PATIENTS WITH ACUTE DESTRUCTIVE PANCREATITIS 4
--

Иванчиков Е., Захаровцев Е., Красавин Е., Шишенинцева Е. MODERN POSSIBILITIES OF DIAGNOSIS AND TREATMENT OF OTOMYCOSES 12
--

A.J. Sebag THE ROLE OF MICROWAVE ABLATION IN THE TREATMENT OF PAPILLARY THYROID CANCER: A SYSTEMATIC REVIEW 18
--

CLINICAL MEDICINE

Багирова Г. RESULTS OF SURGICAL TREATMENT OF HIP INJURIES 22

Надежкин Р., Гаркуш Е., Малиновский Н., Головин А. EFFICIENCY RESULTS OF LAPAROSCOPIC TREATMENT OF LIVER SCHISTOSOCOSIS 26

Балакин Е., Балакин О., Балакин О., Балакин Е., Абрамян А., Голиков А. FACTOR ANALYSIS OF THE RESULTS OF MODERN TREATMENT OF PATIENTS WITH LIVER CIRRHOSIS WITH PORTAL HYPERTENSION 30
--

Ланчук Е., Касимов Г., Шаблыкина А., Багиров Г. HEMORRHAGIC STROKES IN INFANTS 31
--

L. Wang, Huiyan Xie BACTERIAL INFECTION FOLLOWING TOTAL JOINT ARTHROPLASTY 40
--

Ахмедзода Р., Растворова Е., Тод Г., Сабитова А., Зубайдиев С. SURGICAL METHOD FOR THE TREATMENT OF CHRONIC CONDYLITIS IN PATIENTS WITH DOLICHOSIGMA 44

<https://doi.org/10.54500/2790-1203-2023-116-30-34>
ISSN 7629.39+76.29.34
UDC 616-36-004-06:616.149-005.1-089.819

A short communication

THE FACTOR ANALYSIS OF THE RESULTS OF MODERN TREATMENT OF PATIENTS WITH LIVER CIRRHOSIS WITH PORTAL HYPERTENSION

Khakimov M.¹, Matkuliev U.², Batirov D.³, Umarov Z.^{4*}, Allanazarov A.⁵, Rakhimov A.⁶

¹ Head of the department of the 1st faculty and hospital surgery of Tashkent Medical Academy, Tashkent, Uzbekistan.
E-mail: murad_72@mail.ru

² Associate professor of the department of the 1st faculty and hospital surgery of Tashkent Medical Academy, Tashkent, Uzbekistan. E-mail: mat.utkirbek@gmail.com

³ Head of the department of Surgical Diseases and Transplantology of Tashkent Medical Academy Urgench Branch, Urgench, Uzbekistan. E-mail: batirovdavronbek72@gmail.com

⁴ Assistant of the department of Surgical Diseases and Transplantology of Tashkent Medical Academy Urgench Branch. Uzbekistan, 220100, Urgench, st. Al-Khwarizmi, 28. E-mail: doctor_uzz1986@mail.ru

⁵ Associate professor of the department of Surgical Diseases and Transplantology of Tashkent Medical Academy Urgench Branch. Urgench, Uzbekistan. E-mail: allanazarovallanazar1985@gmail.com

⁶ Associate Professor of the department of Surgical Diseases and Transplantology of Tashkent Medical Academy Urgench Branch. Urgench, Uzbekistan. E-mail: arvabek2277@gmail.com

Abstract

The purpose of this study is to study and evaluate the results of conventional methods of endovascular and endoscopic treatment of patients with complicated portal hypertension.

Methods. In our study until 2021, 56 patients were included for analysis. In the course of the research, in 12 cases the endovascular method of percutaneous hepatic embolization of esophagus and gastric varicose vein in combination with splenic artery embolization, endoscopic sclerotherapy method in 23 cases, and endoscopic ligation according to the traditional method in 21 cases were used.

The result of research work has showed that isolated use of endovascular methods (left gastric vein embolization, splenic artery embolization) in patients with liver cirrhosis is characterized by a low 2-year survival rate - 14.2%, with frequent anatomical and technical complications of this procedure - up to 84.7%, unstable primary hemostasis - 40.4%, esophagus and gastric varicose vein are explained by a high rate of recurrence of bleeding from varicose veins - 67.4%, risk of death - 21.8%.

Conclusions. The clinical advantages of EL compared to different methods of ES have been achieved in terms of the rate of ineffective primary hemostasis, the risk of recurrence of bleeding from EGVV, short-term and long-term results, as well as improved 2-year survival rates. However, in general, their isolated use is associated with a relatively low chance of complete prevention of exacerbation of hemorrhagic syndrome relapses, which makes it possible to choose a combined approach to the treatment of patients with LC with PH and risk of bleeding from EGVV requires the development of tactical algorithms.

In endovascular transhepatic embolization of esophageal and gastric varices, good results of treatment in the follow-up period of up to 2 years were very low, while unsatisfactory results of ES prevailed over good results, in turn, the best results were observed in the EL group. The results of endovascular transhepatic embolization in EGVV remain unknown in 1/3 of the studies.

Key words: liver cirrhosis, portal hypertension, endoscopic ligation, endovascular, esophagus and gastric variceal bleeding.

Material received: 18/03/2023

Material accepted for publication: 04/05/2023

Reference for citation: Khakimov M., Matkuliev U., Batirov D., Umarov Z., et al. The factor analysis of the results of modern treatment of patients with liver cirrhosis with portal hypertension. Astana medical journal. 2023; 116: 30-34 . <https://doi.org/10.54500/2790-1203-2023-116-30-34>

Introduction

Globally, the development of surgery in general, and endoscopic surgery in particular, depends on striving to minimize the amount of intervention, regardless of the severity of the patient's clinical condition [1-5]. Evaluating the effectiveness of treatment of patients with portal hypertension (PH) and liver cirrhosis (LC) after endovascular and endoscopic interventions and patient survival are very important tasks [6-8]. Esophagus and Gastric Varicose Vein (EGVV) patients with LC complicated by bleeding, by improving the tactical and technical aspects of various minimally invasive interventions, the aspects of minimally invasive diagnostic and treatment methods remain one of the most relevant today. In this group of patients, the issues of developing treatment-tactical algorithms, as well as evaluating the clinical effect of minimally invasive interventions in the near and long term, remain [9,10].

Currently, worldwide research is ongoing to improve the diagnosis and treatment of bleeding

recurrences due to PH, including: development of new less invasive, less invasive methods of PH correction; Early diagnosis of LC formation, its prognosis and optimal tactics of treatment; Evaluation of treatment efficacy and survival after endovascular and endoscopic interventions in patients with LC with PH [11,12].

Analysis of the data presented in the literature has significantly improved the results of surgical treatment of patients with LC with PH. In this case, the use of minimally invasive endovascular, endoscopic and traditional methods of treatment in turn, in stages, is the main current and determining problem, which means that the need for further improvement of treatment-diagnostic tactics has not lost its relevance[1,10].

The purpose of this study is to study and evaluate the results of conventional methods of endovascular and endoscopic treatment of patients with complicated portal hypertension.

Material and methods

In order to study and evaluate the fundamental analysis of the results of traditional methods of endovascular and endoscopic treatment of patients with complicated PH, studies based on the analysis of the results of treatment of 56 patients with PH and LC were conducted in the department of the 1st Faculty and Hospital Surgery of Tashkent Medical Academy and in the multidisciplinary medical center of Khorezm region in 2018-2021. In the course of the research, in 12 cases the endovascular method of percutaneous hepatic embolization of EGVV in combination with

Splenic Artery Embolization, endoscopic sclerotherapy (ES) method in 23 cases, and endoscopic ligation (EL) according to the traditional method in 21 cases were used. The main criteria for evaluating the effectiveness of endovascular and endoscopic interventions were: the effectiveness of stopping bleeding, the possibility of preventing recurrences of bleeding in the near and long term, the reduction of complications and mortality. Long-term results were evaluated as good, satisfactory and unsatisfactory.

Results

In the near term (hospital phase), the overall effectiveness of the methods in the prevention of recurrence of hemorrhagic syndrome after endovascular transhepatic embolization of EGVV was 66.7% (8 out of 12 patients), 73.9% (17 out of 23) after ES and 90.5% (19 of 21) after EL. It should be noted that anatomical or technical complications were observed in endovascular intervention (54.3%) (which explains such a low level of efficiency), as well as the highest rate of the closest recurrence of bleeding.

Based on the results of the treatment, it is possible to determine the indications for inter-hepatic intervention, which can achieve the best result with minimal risk. The

annotated material shows that in the group of patients with liver failure of class A, the mortality rate during liver intervention was 7.1%, 92.9% of patients were discharged from the hospital with positive dynamics and satisfactory condition.

Thus, isolated use of endovascular methods (left gastric vein embolization, splenic artery embolization) in patients with LC is characterized by a low 2-year survival rate - 14.2%, with frequent anatomical and technical complications of this procedure - up to 64.7%, unstable primary hemostasis - 40.4%, EGVV are explained by a high rate of recurrence of bleeding from varicose veins - 67.4%, risk of death - 21.8%.

Discussion

Indications and prediction of positive results in the use of interhepatic endovascular interventions in this case indicate that the initial state of liver failure in class A is considered. During the development of endoscopic interventions, this method can be considered as an alternative method for esophagus varicose veins, and for gastric varicose veins as a choice method.

The use of EL and ES in acute esophageal-gastric bleeding of portal genesis allows to stop bleeding in high-risk cases, improve the two-year survival of patients, and improve the results compared to patients who underwent interhepatic intervention. In this case, the most important prognostic factors influencing the survival of patients who experienced portal genesis bleeding are their functional class according to Child-Pugh and increased recurrence of bleeding. The main factor associated with increased

recurrence of variceal bleeding after the endoscopic treatment program was the inability to achieve endoscopic eradication of gastric cardial varicose veins, which in our observations was 44.9% and 20.2% after ES and EL, respectively. The use of EL of varicose veins, in comparison with the isolated use of ES, occurs with an increase in the percentage of patients with eradication, which undoubtedly reduces the rate of relapses and deaths. Reducing the risk of recurrence of bleeding from EGVV by carrying out an endoscopic program of treatment allows improving the quality of life of patients with LC.

However, due to the persistence of PH and Liver Failure, the long-term results of the use of endoscopic interventions are not encouraging. This situation indicates

the need to include endovascular interventions aimed at reducing liver failure in the complex of treatment measures.

Conclusions

The clinical advantages of EL compared to different methods of ES have been achieved in terms of the rate of ineffective primary hemostasis, the risk of recurrence of bleeding from EGVV, short-term and long-term results, as well as improved 2-year survival rates. However, in general, their isolated use is associated with a relatively low chance of complete prevention of exacerbation of hemorrhagic syndrome relapses, which makes it possible to choose a combined approach to the treatment of patients with LC with PH and risk of bleeding from EGVV requires the development of tactical algorithms.

In endovascular transhepatic embolization of esophageal and gastric varices, good results of treatment in the follow-up period of up to 2 years were very low, while unsatisfactory results of ES prevailed over good results, in turn, the best results were observed in the EL group. The results of endovascular transhepatic embolization in EGVV remain unknown in 1/3 of the studies.

References

1. Shavkatovich X.M., Ismoilovich M. U., Zaripboyevich U. Z. Portal gipertensiya asoratlarini oldini olish va davolashda kaminzavizli aralashuvlar kompleksining samaradorligi (The effectiveness of a complex of minimally invasive interventions in the prevention and treatment of complications of portal hypertension) [in Uzbek]. Science and innovation, 2022; 1(Special Issue 2): 379-383. [[Google Scholar](#)]
2. Khakimov M. Endovascular Reduction of Splenic Blood Flow in the Treatment of Patients with Liver Cirrhosis. Journal of Advances in Medicine and Medical Research. 2016; 13(5): 1-10. [[Google Scholar](#)]
3. Karimov S., Matkuliev U., Khakimov M. et al. Selection of endoscopic interventions in the prevention of rebleeding in patients with portal hypertension. International Journal of Information Research and Review. 2016; 3: 1832-1836. [[Google Scholar](#)]
4. Каримов Ш.И., Маткулев У.И., Хакимов М.Ш. и др. Оценка эффективности эндоскопических вмешательств у больных с кровотечением из варикозно расширенных вен пищевода и желудка // Вестник ТМА. – 2016. — № 1. – С. 61-66. [[Google Scholar](#)]
- Karimov Sh.I., Matkuliev U.I., Hakimov M.Sh. i dr. Ocena jeffektivnosti jendoskopicheskikh vmeshatel'stv u bol'nyh s krovotecheniem iz varikozno rasshirennyyh ven pishhevoda i zheludka [Evaluation of the effectiveness of endoscopic interventions in patients with bleeding from varicose veins of the esophagus and stomach] [in Russian] Vestnik TMA. 2016; 1; 61-66. [[Google Scholar](#)]
5. Каримов Ш.И., Хакимов М.Ш., Маткулев У.И. Выбор эндоскопических вмешательств в лечении кровотечений из варикозных вен пищевода и желудка // Вахидовские чтения: Республикаанская научно-практическая конференция. - Ташкент, 2014. С. 75-76 [[Google Scholar](#)]
- Karimov Sh.I., Hakimov M.Sh., Matkuliev U.I. Vybor jendoskopicheskikh vmeshatel'stv v lechenii krovotechenij iz varikoznyh ven pishhevoda i zheludka (The choice of endoscopic interventions in the treatment of bleeding from varicose veins of the esophagus and stomach) [in Russian]. Vahidovskie chtenija: Respublikanskaja nauchno-prakticheskaja konferencija. - Tashkent, 2014. -75-76. [[Google Scholar](#)]
6. Маткулев У.И., Умаров З.З. Малоинвазивные вмешательства в профилактике и лечении кровотечений у больных с портальной гипертензией //Научный обозреватель – 2020. - № 10 (118). – С. 47-50
- Matkuliev U.I., Umarov Z.Z. Maloinvazivnye vmeshatel'sta v profilaktike i lechenii krovotechenij u bol'nyh s portal'noj gipertenzijej (Minimally invasive interventions in the prevention and treatment of bleeding in patients with portal hypertension) [in Russian]. Nauchnyj obozrevatel' 2020; 10(118): 47-50.
7. Wen J., Chen X., Wei S., Ma X., Zhao Y. Research progress and treatment status of liver cirrhosis with hypoproteinemia. Evidence-based Complementary and Alternative Medicine, 2022; 24. [[CrossRef](#)]
8. Batirov D.Y., Kh A.A., Rakhitov A. P., Rojobov R. R. Improving the results of simultaneous laparoscopic surgery in patients with chronic deficiency. European journal of modern medicine and practice. 2023; 3(2): 43-45. [[Google Scholar](#)]
9. Хакимов М.Ш., Маткулев У.И., Холматов Ш.Т. Эндоскопический гемостаз при кровотечениях из варикозных вен пищевода и желудка у больных с внепеченочной портальной гипертензией //Материалы научно-практической конференции 21-го международного конгресса ассоциации гепатопанкреатобилиарных хирургов стран СНГ. - Иркутск, 19-21 сентября, 2012 - С. 26-27.
- Hakimov M.Sh., Matkuliev U.I., Holmatov Sh.T. Jendoskopicheskij gemostaz pri krovotechenijah iz varikoznyh ven pishhevoda i zheludka u bol'nyh s vnenepchenochnoj portal'noj gipertenzijej (Endoscopic hemostasis in bleeding from varicose veins of the esophagus and stomach in patients with extrahepatic portal hypertension) [in Russian]. Materialy nauchno-prakticheskoj konferencii 21-go mezhdunarodnogo kongressa associacii hepatopankreatobiliarnykh hirurgov stran SNG. - Irkutsk, 19-21 sentyabrya, 2012. - S. 26-27.
10. Маткулев У.И., Хакимов М.Ш., Асраров А.А. и др. Выбор способа гемостаза при кровотечениях из варикозных вен желудка у пациентов с портальной гипертензией // Актуальные проблемы гепатопанкреатобилиарной хирургии: Материалы 22-го международного конгресса Ассоциации гепатопанкреатобилиарных хирургов стран СНГ. - Ташкент, 7-9 сентября 2015 г. – С. 50
- Matkuliev U.I., Hakimov M. Sh., Asrarov A.A. i dr. Vybor sposoba gemostaza pri krovotechenijah iz varikoznyh ven zheludka u pacientov s portal'noj gipertenzijej (The choice of hemostasis method for bleeding from gastric varicose veins in patients with portal hypertension) [in Russian]. Aktual'nye problemy hepatopankreatobiliarnoj hirurgii: Materialy 22-go mezhdunarodnogo kongressa Associacii hepatopankreatobiliarnykh hirurgov stran SNG. - Tashkent, 7-9 sentyabrya 2015: 50.

11. Маткулиев УИ. Хакимов М.Ш. Ашурев Ш. Э. и др. Прогнозирование повторных кровотечений у больных с портальной гипертензией // Актуальные проблемы гепатопанкреатобилиарной хирургии: Материалы 22-го международного конгресса ассоциации гепатопанкреатобилиарных хирургов стран СНГ. - Ташкент, 7-9 сентября 2015 г. - С. 51 [[Google Scholar](#)]

Matkuliev U.I. Hakimov M.Sh. Ashurov Sh. Je. idr Prognozirovaniye povtornykh krovotечenij u bol'nyh s portal'noj gipertenziiey [Prediction of rebleeding in patients with portal hypertension] [in Russian]. Aktual'nye problemy hepatopankreatobiliarnoj hirurgii: Materialy 22-go mezhdunarodnogo kongressa associacii hepatopankreatobiliarnyh hirurgov stran SNG. - Tashkent, 7-9 sentyabrya 2015 g. - S. 51 [[Google Scholar](#)]

12. Хакимов М.Ш. TIPS как альтернатива открытым портокавальным шuntам // Актуальные проблемы гепатобилиарной хирургии. - 2015. - С. 42-43. [[Google Scholar](#)]

Khakimov M.Sh. TIPS kak al'ternativa otkrytym portokaval'nym shuntam (TIPS as an alternative to open porto-caval shunts) [in Russian]. Aktual'nye problemy hepatobiliarnoj hirurgii. - 2015: 42-43. [[Google Scholar](#)]

ПОРТАЛДЫҚ ГИПЕРТЕНЗИЯМЕН АСҚЫНГАН БАУЫР ЦИРРОЗЫМЕН АУЫРАТЫН НАУҚАСТАРДЫ ҚАЗІРГІ ЗАМАНҒЫ ЕМДЕЛУ НӘТИЖЕЛЕРІНІҢ ФАКТОРЛЫҚ ТАЛДАУЫ

Хакимов М.Ш.¹, Маткулиев О.И.², Батиров Д.Ю.³, Умаров З.З.⁴, Алланазаров А.Х.⁵, Рахимов А.П.⁶

¹ Ташкент медицина академиясының 1-ші факультет және госпитальдық хирургия кафедрасының мензверушісі, Ташкент, Өзбекстан. E-mail: murad_72@mail.ru

² Ташкент медицина академиясының 1-факультет және госпитальдық хирургия кафедрасының доценті Ташкент, Өзбекстан. E-mail: mat.utkirbek@gmail.com

³ Ташкент медицина академиясы Уралниш филиалының хирургиялық аурулар және трансплантология кафедрасының мензверушісі, Уралниш, Өзбекстан. E-mail: batirovdaurovbek72@gmail.com

⁴ Ташкент медицина академиясы Уралниш филиалының хирургиялық аурулар және трансплантология кафедрасының асистенті, Уралниш, Өзбекстан. E-mail: doctor_uzz1986@mail.ru

⁵ Ташкент медицина академиясы Уралниш филиалының хирургиялық аурулар және трансплантология кафедрасының доценті Уралниш, Өзбекстан. E-mail: allanazarovallanazar1985@gmail.com

⁶ Ташкент медицина академиясы Уралниш филиалының хирургиялық аурулар және трансплантология кафедрасының доценті Уралниш, Өзбекстан. E-mail: anvarbek2277@gmail.com

Түйіндеме

Зерттеудің мақсаты - асқынған порталдық гипертензиясы бар науқастарды эндоваскулярық және эндоскопиялық емдеудің дәстүрлі әдістерінің нәтижелерін зерттеу және бағалау болды.

Әдістері. 2021 жылға дейінде зерттеуіміз таңдауға 56 пациентті қамтыйды. Зерттеу барысында өнеш пен асқазанның варикозды венапарының төрі арқылы бауыр эмболизациясының эндоваскулярық адісі 12 жағдайда көкбауыр артериясының эмболизациясы, эндоскопиялық склеротерапия 23 жағдайда және 21 жағдайда дәстүрлі әдіс бойынша эндоскопиялық байлау қолданылды.

Зерттеу нәтижелері көрсеткендегі, бауыр циррозы бар науқастарда эндоваскулярық әдістерді (асқазанның сол жақ венасының эмболизациясы, көкбауыр артериясының эмболизациясы) оқшауланған қолдану 2 жылдық емір сурудің тәмем деңгейімен 14,2%, жиі анатомиялық және осы операцияның техникалық асқынұлары. операциялар - 64,7% дейін, тұрақсыз бірнешілік асемтас - 40,4%, өнеш пен асқазанның варикозды венапары варикозды тамырлардан қан кетудің қайталану жиілігімен туындылады - 67,4%, алғы қаупі - 21,8%.

Қорытынды. Тиімді бірнешілік гемостаздың жиілігі, варикозды көнегең венаның эмболиясынан соң қайталананатын қан кету қаупі, жедел және ұзақ мерзімді нәтижелер және 2 жылдық емір суруда көрсеткіштердің жақсаруы бойынша ЭС өртүрлі әдістермен салыстырғанда ЭЛ клиникалық артықшылықтарына қол жеткізілді. Алайда, тұмастай алғанда, оларды оқшауланған қолдану геморрагиялық синдромның қайталануының толық алдың алудың салыстырмалы түрде тәмем ықтималдығымен байланысты, бұл портальды гипертензиясы бар науқастарды емдеуге біріктірілген тесілді таңдауға мүмкінді береді және варикозды көнегең венаның эмболиясынан соң қан кету қаупін жоятын тәжірибелі алгоритмдерді құрастыру.

Өнеш пен асқазанның варикозды көнегеңінің эндоваскулярық трансбауыр эмболизациясында 2 жылға дейінде бағылау кезеңінде жақсы емдеу нәтижелері етте тәмем болды, ал ЭС қанагаттанарлықсыз нәтижелері жақсыдан басым болды, ал көзегінде өн жақсы нәтижелер ЭЛ тобында. Байқалды. Эндоваскулярық трансбауырлық эмболизациясының нәтижелері зерттеулердің 1/3 белгінде белгісіз болып қалады.

Түйін сөздер: бауыр циррозы, порталдық гипертензия, эндоскопиялық байлау, варикозды веналардың эндоваскулярық эмболиясы, внештеп және асқазаннан қан кету.

ФАКТОРНЫЙ АНАЛИЗ РЕЗУЛЬТАТОВ СОВРЕМЕННОГО ЛЕЧЕНИЯ БОЛЬНЫХ ЦИРРОЗОМ ПЕЧЕНИ ОСЛОЖНЕННЫМ ПОРТАЛЬНОЙ ГИПЕРТЕНЗИЕЙ

Хакимов М.¹, Маткулиев У.², Батиров Д.³, Умаров З.^{4*}, Алланазаров А.⁵, Рахимов А.⁶

¹ Заведующий кафедрой 1-го факультета и госпитальной хирургии Ташкентской медицинской академии.

E-mail: murad_72@mail.ru

² Доцент кафедры 1-го факультета и госпитальной хирургии Ташкентской медицинской академии, Ташкент,

Узбекистан. E-mail: mat.utkirbek@gmail.com

³ Заведующий кафедрой хирургических болезней и трансплантологии Ургенчского филиала Ташкентской медицинской академии, Ургенч, Узбекистан. E-mail: batirovdavronbek72@gmail.com

⁴ Ассистент кафедры хирургических болезней и трансплантологии Ургенчского филиала Ташкентской медицинской академии, Ургенч, Узбекистан. E-mail: doctor_uzz1986@mail.ru

⁵ Доцент кафедры хирургических болезней и трансплантологии Ургенчского филиала Ташкентской медицинской академии, Ургенч, Узбекистан. E-mail: allanazarovallanazar1985@gmail.com

⁶ Доцент кафедры хирургических болезней и трансплантологии Ургенчского филиала Ташкентской медицинской академии, Ургенч, Узбекистан. E-mail: anvarbek2277@gmail.com

Резюме

Цель исследования - изучить и оценить результаты общепринятых методов эндоваскулярного и эндоскопического лечения больных с осложненной портальной гипертензией.

Материал и методы. В наше исследование до 2021 года для анализа были включены 56 пациентов. В ходе исследования в 12 случаях использовали эндоваскулярный метод чрескожной печеночной эмболизации варикозно расширенных вен пищевода и желудка в сочетании с эмболизацией селезеночной артерии, метод эндоскопической склеротерапии в 23 случаях и эндоскопическое лигирование по традиционной методике в 21 случае.

Результаты исследования показали, что изолированное применение эндоваскулярных методов (эмболизация левой желудочной вены, эмболизация селезеночной артерии) у больных циррозом печени характеризуется низкой 2-летней выживаемостью - 14,2%, с частыми анатомо-техническими осложнениями этой операции. операции - до 64,7%, нестабильны первичный гемостаз - 40,4%, варикозное расширение вен пищевода и желудка объясняются высокой частотой рецидивов кровотечений из варикозно расширенных вен - 67,4%, риском летального исхода - 21,8%.

Выводы. Достигнуты клинические преимущества ЭЛ по сравнению с различными методами ЭС по частоте незэффективного первичного гемостаза, риска рецидива кровотечения из ЭГВВ, непосредственных и отдаленных результатов, а также улучшены 2-х годовых показатели выживаемости. Однако в целом их изолированное применение связано с относительно низкой вероятностью полной профилактики обострения рецидивов геморрагического синдрома, что позволяет выбрать комбинированный подход к лечению больных РЛ с ЛГ, а риск кровотечения из ЭГВВ требует разработка тактических алгоритмов.

При эндоваскулярной чрескожной эмболизации варикозно расширенных вен пищевода и желудка хорошие результаты лечения в сроки наблюдения до 2 лет были очень низкими, при этом неудовлетворительные результаты ЭС преобладали над хорошими, в свою очередь, лучшие результаты наблюдались при группе ЭЛ. Результаты эндоваскулярной чрескожной эмболизации при ЭГВВ остаются неизвестными в 1/3 исследований.

Ключевые слова: цирроз печени, портальная гипертензия, эндоскопическое лигирование, эндоваскулярное, пищеводное и желудочное кровотечение из варикозно расширенных вен.