



Tashkent Medical
Academy



Siberian state medical
university

International scientific and practical conference of young scientists

“ISSUES OF BIOPHYSICS IN MEDICINE”



ISSN 2181-7812

11 May, 2023

MINISTRY OF HIGHER AND SECONDARY SPECIAL EDUCATION
OF THE REPUBLIC OF UZBEKISTAN

MINISTRY OF HEALTHCARE
TASHKENT MEDICAL ACADEMY

**ABSTRACT BOOK OF THE INTERNATIONAL SCIENTIFIC AND PRACTICAL
CONFERENCE OF YOUNG SCIENTISTS “ISSUES OF BIOPHYSICS IN MEDICINE”**

TASHKENT-2023

O'ZBEKİSTON RESPUBLİKASI
OLIY VA O'RTA MAXSUS TA'LIM VAZIRLIGI

O'ZBEKİSTON RESPUBLİKASI SOĞ'LIQNI SAQLASH VAZIRLIGI

TOSHKENT TIBBIYOT AKADEMIYASI

**“TIBBIYOTDA BIOFİZİKA MASALALARI” MAVZUSIDAGI
YOSH OLIMLARNING XALQARO İLMİY-AMALİY KONFERENSIYA TO'PLAMI**

TOSHKENT-2023

МИНИСТЕРСТВО ВЫСШЕГО И СРЕДНЕГО СПЕЦИАЛЬНОГО
ОБРАЗОВАНИЯ РЕСПУБЛИКИ УЗБЕКИСТАН

МИНИСТЕРСТВО ЗДРАВООХРАНЕНИЯ РЕСПУБЛИКИ
УЗБЕКИСТАН

ТАШКЕНТСКАЯ МЕДИЦИНСКАЯ АКАДЕМИЯ

**СБОРНИК МАТЕРИАЛОВ МЕЖДУНАРОДНОЙ НАУЧНО-ПРАКТИЧЕСКОЙ
КОНФЕРЕНЦИИ МОЛОДЫХ УЧЁНЫХ “ВОПРОСЫ БИОФИЗИКИ В
МЕДИЦИНЕ”**

Ташкент – 2023

CHIEF EDITOR
professor Shadmanov A.K.

EDITORIAL TEAM:

Gusakova S.V. – Head of Biophysics and Functional Diagnostics Division of Siberian State Medical University (SibMed), professor	Aliyev S.U. – Dean of the Department Pharmacy, management, medical biology, medical bioengineering and higher nursing faculty of Tashkent Medical Academy (TMA), Associate professor
Bazarbayev M.I. head of the "Biomedical Engineering, Informatics and Biophysics" department of Tashkent Medical Academy (TMA), Associate professor	Mullojonov I. – Associate professor of the department "Biomedical Engineering, Informatics and Biophysics" of TMA
Maxsudov V.G. – senior lecturer of the department "Biomedical Engineering, Informatics and Biophysics" of TMA, PhD	Ermetov E.Ya. – senior lecturer of the department "Biomedical Engineering, Informatics and Biophysics" of TMA
Sobirjonov A.Z. – senior lecturer of the department "Biomedical Engineering, Informatics and Biophysics" of TMA	Raximov B.T. – assistant of the department "Biomedical Engineering, Informatics and Biophysics" of TMA
Abdujabbarova U.M. – assistant of the department "Biomedical Engineering, Informatics and Biophysics" of TMA	Sayfullayeva D.I. – assistant of the department "Biomedical Engineering, Informatics and Biophysics" of TMA

ГЛАВНЫЙ РЕДАКТОР
проф. Шадманов А.К.

РЕДАКЦИОННАЯ КОЛЛЕГИЯ:

Гусакова С.В. – доктор медицинских наук, заведующая кафедрой Биофизики и функциональной диагностики Сибирского государственного медицинского университета (СибГМУ), профессор	Алиев С.У. – Декан факультета “Фармация, менеджмент, медицинская биология, биомедицинская инженерия и ВСД Ташкентского медицинского академии (ТМА), доцент
Базарбаев М.И. – заведующий кафедрой «Биомедицинского инженерии, информатики и биофизики» ТМА, доцент	Муллоюнов И. – доцент кафедрой «Биомедицинского инженерии, информатики и биофизики» ТМА
Максудов В.Г. – старший преподаватель кафедры «Биомедицинского инженерии, информатики и биофизики» ТМА, PhD	Эрметов Э.Я. – старший преподаватель кафедры «Биомедицинского инженерии, информатики и биофизики» ТМА
Собиржонов А.З. – старший преподаватель кафедры «Биомедицинского инженерии, информатики и биофизики» ТМА	Рахимов Б.Т. – ассистент кафедры «Биомедицинского инженерии, информатики и биофизики» ТМА
Абдузаббарова У.М. – ассистент кафедры «Биомедицинского инженерии, информатики и биофизики» ТМА	Сайфуллаева Д.И. – ассистент кафедры «Биомедицинского инженерии, информатики и биофизики» ТМА

BOSH MUHARRIR
professor Shadmanov A.K.

TAHRIR HAYATI

Gusakova S.V. – tibbiyot fanlari doktori, Sibir davlat tibbiyot universitetining Biofizika va funksional diagnostika kafedrasи mudiri, professor	Aliyev S.U. – TTA, Farmatsiya, menejment, tibbiy biologiya, tibbiy biomuhandislik, oly ma'lumotli hamshira fakulteti dekanı, dotsent
Bazarbayev M.I. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи mudiri, dotsent	Mullojonov I. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи dotsenti
Maxsudov V.G. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи katta o'qituvchisi, PhD	Ermetov E.Ya. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи katta o'qituvchisi
Sobirjonov A.Z. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи katta o'qituvchisi	Raximov B.T. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи assistenti
Abdujabbarova U.M. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи assistenti	Sayfullayeva D.I. – TTA Biotibbiyot muhandisligi, informatika va biofizika kafedrasи assistenti

PERINATAL HYPOXIA EFFECTS ON THE ENERGETIC FUNCTION OF BRAIN NEURONAL MITOCHONDRIA

Yeliubayev K.O.*, Kairat B.K.

Al-Farabi Kazakh National University, Almaty, Kazakhstan

*E-mail: kanat.yeliubayev@gmail.com

Abstract. This article presents the biophysical and molecular mechanisms of the response of brain neurons to hypoxia, describing disturbances in mitochondrial energy functions and ion homeostasis in cells exposed to hypoxia and ischaemia.

Keywords: perinatal hypoxia, ischaemia, brain, neuron, cell homeostasis, necrosis, apoptosis.

Perinatal hypoxia is an important factor in the development of the central nervous system. The more advanced neonatal intensive care technologies improve survival rates, but do not prevent neurological disorders, resulting in increased morbidity in the adult population [1-3].

Perinatal hypoxia is a pathological process associated with oxygen deprivation during pregnancy and the early neonatal period. Perinatal hypoxia decreases or stops the supply of oxygen to the body, and under-oxidised products of metabolism accumulate in the blood. The cerebral ischaemia (decreased cerebral blood flow) is an equally important factor in the development of neuronal abnormalities in the perinatal period. In this condition there is not only oxygen deprivation, but also a reduction in glucose supply to the brain tissue [4]. The majority of neurological abnormalities caused by perinatal hypoxia result from a combination of hypoxia and ischaemia [5]. The placenta dysfunction, prolonged labour, premature delivery and cardio-respiratory disorders of the mother cause perinatal fetal hypoxia [6]. The intrauterine hypoxia and asphyxia of the newborn is a major pathogenetic factor in brain damage. The prenatal hypoxia rate is 1-6 per 1 000 infants and is one of the main causes of perinatal mortality [4]. The high incidence and very severe consequences for the child make research into the neurochemical mechanisms underlying the effects of perinatal hypoxia urgent [2].

The developing brain is very sensitive to hypoxic damage as its oxygen requirements are very high compared to other organs and tissues [7]. Poor oxygen supply (hypoxia) and/or reduced blood flow (ischaemia) activate various cytotoxic agents and cell death pathways leading to brain damage [8]. The neuropathological processes are triggered during the acute phase of hypoxia/ischaemia and continue during reperfusion [9]. There are various pathogenic mechanisms underlying hypoxic

damage, such as accumulation of toxic metabolites including glutamate, impaired Ca^{2+} exchange, release of pro-inflammatory mediators, accumulation of iron, overproduction of free radicals, reduction of energy stores and synthesis of macro-ergic compounds [10]. The reduction of macroergs and increased levels of intracellular and extracellular damaging factors lead to disruption of cellular homeostasis [9].

The activity and excitability of neurons is directly regulated in response to O_2 reduction by altering membrane channel conduction. Plasma membrane of brain neurons contains a specific type of potassium channels whose subunits form four transmembrane groupings and two transmembrane pores (background K2P channels) [11]. Their function is related to maintaining cell membrane potential at rest, as well as to the regulation of K^+ homeostasis and neuronal excitability. Some hormones and neurotransmitters, anaesthetics as well as physico-chemical factors such as hypoxia and changes in extra- and intracellular pH modulate the effects of background K2P channels. These K2P channels are potential-independent and permanently active. Their blockade causes depolarisation and their activation causes hyperpolarisation of the neuronal membrane. The family K2P channels of mammals are subdivided into 6 subfamilies. TASK-1/TASK-3 types of K2P channels are sensitive to hypoxia and significantly affect neuronal excitability. Hypoxia and acidosis cause TASK-1/TASK-3 channels to be blocked, leading to membrane depolarisation and neuronal activation [11].

Hypoxia-induced brain damage develops through processes that occur during the acute phase of hypoxia-ischaemia and continue into the reperfusion phase. Oxygen and glucose deprivation in the background of hypoxia-ischaemia or immediately afterwards causes primary energy deficiency [12]. The reduction in oxygen availability leads to impaired electron transport in the mitochondrial respiratory enzyme chain, with the result that the associated process of oxidative phosphorylation suffers, which reduces the synthesis of macroergic compounds such as ATP. There is an energy deficit in the cells. Glucose metabolism is then switched mainly to the anaerobic pathway, leading to the accumulation of lactate and the development of acidosis. Lactate accumulation plays a positive role in adapting to O_2 deficiency, but with increasing levels lactate has a damaging effect [11].

Additionally, disruption of ATP synthesis leads to impairment of energy-dependent homeostatic functions, such as maintenance of cell membrane potential. The ATP-dependent Na^+/K^+ pump is disrupted, leading to intracellular accumulation

of Na^+ , Ca^{2+} ions and water, and cytotoxic oedema develops. As a result, cell death occurs along the necrotic pathway [13]. The depolarisation of the membranes of neurons induces the release of glutamate, leading to further cellular excitation and excessive Ca^{2+} input into the cytoplasm. High Ca^{2+} concentration leads to NO-synthase activation (increased NO content), formation of free radicals, activation of phospholipase (membrane lipid degradation), proteases (protein degradation) and nucleases (DNA degradation). The effect of intracellular Ca^{2+} current on the endothelium of capillaries and arterioles contributes to vasoconstriction and worsens ischaemia. The subsequent reperfusion aggravates the metabolic abnormalities in the brain. Especially, tissue re-oxygenation after an ischaemic episode increases the amount of free oxygen in the cells, which is predominantly used through the oxygenase pathway, being converted into reactive oxygen species (the so-called oxygen paradox). In this period, the damage to cell structure and function is often more severe than in ischaemia [11].

Nonetheless, the development of neuronal damage is not limited to the period of hypoxia/reperfusion. The animal experiments showed significant recovery of cellular metabolism after the end of neonatal hypoxic exposure, followed by a secondary energy deficit [14]. Generally, 6-48 hours after acute hypoxia, cellular energy metabolism is again impaired due to an accumulation of mitochondrial damage, leading to the release of cytotoxic enzymes and pro-apoptotic proteins from the mitochondria [15]. Consequently, some of the neurons that survived the initial energy deficiency die after hours or days mainly through the apoptotic pathway [16].

In summary, despite numerous studies, the long-term effects of perinatal hypoxia and their underlying mechanisms are poorly understood to date. The study of the consequences of perinatal hypoxia and the search for safe and effective methods of their treatment remain the most important task of modern medicine and neuroscience.

References

- 1.Bratek E., Ziembowicz A., Salinska E. Pretreatment with group II metabotropic glutamate receptor agonist LY379268 protects neonatal rat brains from oxidative stress in an experimental model of birth asphyxia // Brain Sci. - 2018. - Vol. 8, № 3. - P. 1-11.
- 2.Millar L.J., Shi L., Hoerder-Suabedissen A., Molnár Z. Neonatal hypoxia ischaemia: mechanisms, models, and therapeutic challenges // Front. Cell. Neurosci. -

2017. - Vol. 11. - P. 1-36.

3.Riljak V., Kraf J., Daryanani A., Jiruška P., Otáhal J. Pathophysiology of perinatal hypoxic- ischemic encephalopathy - biomarkers, animal models and treatment perspectives. // Physiol. Res. - 2016. - Vol. 65, Suppl. 5. - S533-S545.

4.Kurinczuk J.J., White-Koning M., Badawi N. Epidemiology of neonatal encephalopathy and hypoxic-ischaemic encephalopathy // Early Hum. Dev. - 2010. - Vol. 86, № 6. - P. 329-338.

5.Hossain M.A. Molecular mediators of hypoxic-ischemic injury and implications for epilepsy in the developing brain // Epilepsy Behav. - 2005. - Vol. 7, № 2. - P. 204-213.

6.Golan H., Huleihel M. The effect of prenatal hypoxia on brain development: Short- and long- term consequences demonstrated in rodent models // Dev. Sci. - 2006. - Vol. 9, № 4. - P. 338-349.

7.Wang Z., Zhan J., Wang X., Gu J., Xie K., Zhang Q., Liu D. Sodium hydrosulfide prevents hypoxia-induced behavioral impairment in neonatal mice // Brain Res. - 2013. - Vol. 1538. - P. 126-134.

8.Hossain M.A. Hypoxic-ischemic injury in neonatal brain: involvement of a novel neuronal molecule in neuronal cell death and potential target for neuroprotection // Int. J. Dev. Neurosci. - 2008. - Vol. 26, № 1. - P. 93-101.

9.Chen A.I., Xiong L.-J., Tong Y., Mao M. The neuroprotective roles of BDNF in hypoxic ischemic brain injury // Biomed. Reports. - 2013. - Vol. 1, № 2. - P. 167-176.

10.Amato M., Donati F. Update on perinatal hypoxic insult: mechanism, diagnosis and interventions // Eur. J. Paediatr. Neurol. - 2000. - Vol. 4, № 5. - P. 203-209.

11.Lesage F. Pharmacology of neuronal background potassium channels // Neuropharmacology. - 2003. - Vol. 44, № 1. - P. 1-7.

12.Lai M.-C., Yang S.-N. Perinatal hypoxic-ischemic encephalopathy. // J. Biomed. Biotechnol. - 2011. - Vol. 2011. - ID 609813.

13.Iwata O., Iwata S. Filling the evidence gap: how can we improve the outcome of neonatal encephalopathy in the next 10 years? // Brain Dev. - 2011. - Vol. 33, № 3. - P. 221-228.

14.Hagberg H., Mallard C., Ferriero D.M. & et al. The role of inflammation in perinatal brain injury // Nat. Rev. Neurol. - 2015. - Vol. 11, № 4. - P. 192-208.

15.Ten V.S., Starkov A. Hypoxic-ischemic injury in the developing brain: The

role of reactive oxygen species originating in mitochondria // Neurol. Res. Int. - 2012. - Vol. 2012. - ID 542976.

16.Hassell K.J., Ezzati M., & et al. New horizons for newborn brain protection: enhancing endogenous neuroprotection // Arch. Dis. Child. - 2015. - Vol. 100, № 6. - F541-F552.

17.E.Ya.Ermetov M.I.Bazarbayev,U.A.Bozarov, V.G.Maxsudov. Application of differential equations in the field of medicine. International Journal of Engineering Mathematics. 2023. Vol5, #1, -P.7-14.

18.V.G. Maxsudov, E.Ya. Ermetov, Z. R. Jo‘rayeva. Types of physical education and the technologies of organization of matters in the modern education system. Fan, ta’lim va amaliyot integratsiyasi 2022. Vol.4. –P29-34.

19.Эрметов Э.Я. Максудов В.Г. Использование кейс-метода в процессе формирования креативной компетенции будущего учителя математики. Материалы VII учебно-методической конференции "Современные высшее медицинское образование: проблемы, зарубежный опыт, перспективы". 2022/2/25. С.51-52

20.Б Рахимов. The role of innovative educational technologies in teaching biophysics. research and education. 2023. issn: 2181-3191 volume 2 | issue 3 | 202 91-99.

21.Б Рахимов, Х Мухитдинов, З Жўраева. Алгоритм обучения биофизике с использованием инновационных образовательных технологий. 30.03.2023 Innovative Development in Educational Activities issn: 2181-3523 volume 2 issue 6 2023. 191-200.

22.Базарбаев М И., Сайфуллаева Д И., Рахимов Б Т., Жўраева З Р. Роль информационных технологий в медицине и биомедицинской инженерии в подготовке будущих специалистов в период цифровой трансформации в образовании. 10.10.2022. Т.Т.А. Ахборотномаси. 8-13.

23.B.T. Raximov. Современное состояние биофизики и особенности преподавания биофизики в медицинском вузе. Formation of psychology and pedagogy as interdisciplinary sciences. Italia © Sp. z o. o. "CAN", 2021 © Authors, 18-27.

24.Рахимов Б.Т.,Базарбаев М.И.,Собиржонов А.З., Состояние проблемы подготовки студентов-медиков к решению профессиональных задач в обучении биофизике. New Day in Meditcina. www.bsmi.uz <https://newdaymedicine.com> E: ndmuz@mail.ru. 4/54/200-207

TABLE OF CONTENTS
MUNDARIJA
ОГЛАВЛЕНИЕ

SECTION №1 ISSUES OF BIOPHYSICS IN MODERN MEDICINE СЕКЦИЯ №1. ВОПРОСЫ БИОФИЗИКИ В СОВРЕМЕННОЙ МЕДИЦИНЕ	4
THE ROLE OF NA ⁺ ,K ⁺ -ATPASE IN THE MECHANISMS OF VOLUME- DEPENDENT REGULATION OF CONTRACTORY ACTIVITY OF RAT PULMONARY ARTERY SMOOTH MUSCLE CELLS IN HYPOXIA <i>Koshuba S.O.,</i> <i>Prshemysky M.A., Rashkauskaite V.A., Golovanov E.A.</i>	4
PERINATAL HYPOXIA EFFECTS ON THE ENERGETIC FUNCTION OF BRAIN NEURONAL MITOCHONDRIA <i>Yeliubayev K.O.[*], Kairat B.K.</i>	6
БИОФИЗИКА И ЕЕ ЗНАЧЕНИЕ В СФЕРЕ МЕДИЦИНСКОГО ОБРАЗОВАНИЯ <i>Собиржонов А.З.</i>	11
ИССЛЕДОВАНИЕ АЧТВ С ПОМОЩЬЮ КОАГУЛОМЕТРА HUMACLOT JUNIOR <i>Курбонова З.Ч.¹, Полванхонов С.Н.², Имамов Э.З.², Назиров К.Х.²,</i> <i>Абсалямова И.И.²</i>	13
TIBBIYOTDA QO'LLANILAYOTGAN YANGI ASBOB USKUNALAR VA ULARDA METAMATIKA VA FIZIKA FANLARINING AHAMIYATI. <i>Tuxtaxodjayeva F.Sh., Murodullayev M.N.</i>	17
ACCURACY AND RELIABILITY OF MEDICAL DEVICES EQUIPMENT <i>Abdurazzokov J.T., Mamadalieva U.P., Abdullaeva N.U.</i>	23
РОЛЬ ПОТЕНЦИАЛ-ЗАВИСИМЫХ КАЛЬЦИЕВЫХ КАНАЛОВ Т-ТИПА В ГЕНЕРАЦИИ СИНХРОННОЙ АКТИВНОСТИ НЕЙРОНОВ ГИППОКАМПА ПРИ ГИПЕРВОЗБУЖДЕНИИ <i>Сейткадыр К.А., Тулеуханов С.Т., Зинченко В.П.,</i> <i>Кайрат Б.К.</i>	24
TALABALARNI TIBBIYOT OLIY TA'LIM MUASSASALARIDA BIOFIZIKA FANIDAN LABORATORIYA ISHLARINI O'TKAZISH UCHUN TAYYORLASH VA LABORATORIYA MASHG'ULOTLARINI O'TKAZISH METODIKASI <i>Maxkamova M.B., Akromova M.A.</i>	30
GEMATOLOGIK ANALIZATORLARNING ISHLASH PRINSIPI Курбонова З.Ч. ¹ , Полванхонов С.Н. ² , Имамов Э.З. ² , Назиров К.Х. ² , Абсалямова И.И. ²	34
TIBBIYOT OLIY TA'LIM MUASSASALARIDA BIOFIZIKA DARSALARIDA KASBIY MAZMUNDORLIKNI IFODALOVCHI SIFAT MASALALARINI TUZISH, TANLASH VA ULARNI YECHISH METODIKASI Maxkamova M.B... 37	